

First Survey of Epidemiology and Associated Risk Factors of Human Intestinal Parasitic Infections among Gonabad Residents, Northeast of Iran

Mitra Salehi¹, Saber Razavi¹, Nasim khajavian², Hamed Ghasemzadeh-moghaddam¹, Nima Firouzeh^{1*}, Mohammad Hassan Minooeian Haghghi^{3*}

ABSTRACT

OBJECTIVE: To investigate the prevalence of parasitic infections in the human population and identify the related risk factors of Gonabad city, southeast of Khorasan Razavi province, northeastern Iran.

METHODOLOGY: This cross-sectional study was performed from March to April 2019 in Gonabad city in Khorasan Razavi province in northeastern Iran. One thousand thirty-eight stool samples were examined using parasitological methods like direct wet-mounting, formalin-ether concentration and Gomori's trichrome staining. Moreover, the socio-demographic data of inhabitants were recorded. All statistical analyses were performed using SPSS software (version 16). Qualitative variables were reported as numbers and percentages.

RESULTS: The overall intestinal parasitic infection prevalence was 18.1 %, whereas this value for protozoan agents was 16.7% (95% CI 14.14 to 19) and 1.3% (95% CI 0.7 to 2) for helminth parasites. The highest parasitic agent prevalence belongs to *Blastocystis sp* (12.4 %), *Entamoeba coli* (1.8%) and *Enterobius vermicularis* (1.2%), respectively. *Giardia lamblia* (1.1%), *Entamoeba histolytica /dispar* (0.3%), *Iodamoeba butschlii* (0.3%), *Chilomastix mesnili* (0.3%), *Hymenolepis nana* (0.2 %) *Dientamoeba fragilis* (0.2%) and *Trichomonas hominis* (0.3%) were other identified parasites. Multivariate logistic regression analysis revealed a significant association of parasitic intestinal infections with factors such as age, residency status and source of drinking water (P<0.05).

CONCLUSION: Our findings approved that the protozoan agents have a relatively high prevalence among the studied population. So, it could be severe enough to alarm the entities implementing healthcare services measures for parasitic intestinal infection control.

KEYWORDS: Intestinal parasites, helminth, Prevalence, Iran, protozoan

INTRODUCTION

Despite rapid progress in healthcare services performance and public hygiene status in developing countries, parasitic intestinal infection remains a significant public health concern. Annually, it is evaluated that more than three billion cases worldwide experience intestinal parasite infections, whereas 450 million of this population exhibit clinical manifestations. The annual mortality rate of parasitic intestinal infection is estimated to be nearly 16 million. Parasitic intestinal infections are commonly present with clinical

symptoms such as weight loss, malnutrition, abdominal pain or cramping, diarrhoea, vomiting and anemia. These symptoms are particularly prevalent in children and older adults. Additionally, more dangerous complications such as intestinal obstruction, myocarditis, cholecystitis and appendicitis were reported¹¹.

In Iran, the intestinal parasite's prevalence varies between 4.7 and 56% in the healthy human population²². There were many reports of parasitic intestinal infections from different parts of Iran, such as *Blastocystis sp* and *Giardia lamblia*, among Iran's most common pathogenic protozoa. The oral faecal and contaminated food/water are intestinal parasitic agents' most common transmission routes²⁴. Intestinal parasitic infections caused by *Giardia lamblia*, *Blastocystis*, *Entamoeba* and *Cryptosporidium spp(s)* are the most important causes of diarrhea in developing countries²⁴. Over the past four decades, approximately 50% of the Iranian population has experienced pathogenic intestinal parasite infections³¹.

However, several efforts have been made in Iran to decrease the parasitic burden, but the determination

¹Vector-Borne Disease Research Center, North Khorasan University of Medical Sciences, Bojnurd, Iran

²Department of Epidemiology and Biostatistics, School of Health, Social Determinants of Health Research Center, Gonabad University of Medical Sciences, Gonabad, Iran

³Department of Medical Microbiology, Faculty of Medicine, Gonabad University of Medical Sciences, Gonabad, Iran

*Correspondence: nimafrouzeh4@gmail.com, drminooeian@gums.ac.ir

doi: 10.22442/jlumhs.2024.01093

Received: 22-11-2023

Revised: 03-02-2024

Accepted: 22-02-2024

Published Online: 14-05-2024



of intestinal parasite distribution/incidence pattern remains a significant health problem. The prevalence of parasitic intestinal infection in any region has been proven to depend on socioeconomic, personal and public hygiene, and geographic educational status¹². So, it is urgent to identify these factors and establish an efficient prevention/control program for parasitic infections in each zone separately.

Based on our knowledge, there is no scientific report about the distribution of parasitic intestinal infections and associated risk factors in Gonabad city in Khorasan Razavi province in northeastern Iran. Therefore, the present study was designed for the first time to determine the prevalence and associated risk factors with parasitic intestinal infections in the studied human population.

METHODOLOGY

Sample size


Based on the prevalence of infections in the recent study ($p=0.17$), the estimated sample size was calculated using the formula below. Considering a 95% confidence interval and 80% test power, 957 people were selected, which was increased to 1038 individuals, assuming a 10% attrition rate (Abbaszadeh Afshar et al., 2020).

$$n = \frac{(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta})^2 * p(1-p)}{d^2} = \frac{(1.96 + 0.84)^2 * (0.17 * 0.83)}{(0.2 * 0.17)^2} = 957$$

In this research, stratified random sampling was performed. Thus, comprehensive health service centres in Gonabad were considered classes; then, according to the number of people covered by each centre, the samples were randomly selected.

Study area

This cross-sectional study was performed from March to April 2019 in Gonabad city in Khorasan Razavi province in northeastern Iran (34°21'10"N 58°41'01" E). It has an area of about 2188.91 km² and over 36367 people. The city is located in a semi-arid area, with an elevation between 894 and 2774 meters above sea level and has hot summers and cool winters. Gonabad weather station data shows its rainfall and average annual temperature are 149–155 mm/year and 16.4–17.3°C, respectively. This city is famous for its unique groundwater supply, named Qanats. It was added to UNESCO's list of World Heritage Sites in 2007 due to its unique features, like historical sites and ancient artefacts²³. **Figure 1** shows the geographical location of Iran and Gonabad.



Sample collection and laboratory analysis

In the first step, written informed consent was obtained from all of our study's volunteers. Next, they filled out a structured checklist containing individual socio-demographic and health data like sex, age, occupation, education, drinking water source, raw vegetable consumption and family size (Small = less than 5, Large = more than 5). Signing informed consent, delivering stool specimens and not receiving anti-parasitic treatments in the last 21 days or during data collection were our inclusion criteria in the present examination.

Here, 1038 stool samples were collected from Gonabad inhabitants. Stool specimens were collected in sterile plastic containers with numbered labels and promptly transferred to the Department of Parasitology and Mycology, Gonabad University of Medical Sciences.

All collected samples were initially examined through macroscopic observation regarding consistency, the existence of blood or mucus, color and adult forms of helminth parasites. For watery/bloody stool samples, fresh smears were prepared and examined with normal saline (0.85% NaCl solution) and Lugol's iodine solution. All stool samples were stored in buffered formalin (10 %) for further parasitological techniques. According to the previously described protocol, besides wet mount, concentration techniques using formalin-ethyl acetate and Merthiolate-iodine-formaldehyde solution (MIF) were applied²⁰. The same expert observed the prepared slides (blinded to participants' information) through different microscope magnifications. If parasite forms like eggs, cysts, larvae, and/or trophozoites were seen in at least one of the three techniques, that sample would be considered positive, and then the permanent trichrome staining technique would be used to confirm detection.

Statistical analysis

All statistical analyses were performed using SPSS software (version 16). Qualitative variables were reported as numbers and percentages.

The logistic regression model was adopted to determine the factors related to parasitic intestinal infections. Therefore, the variables that had $P < 0.2$ in simple logistic regression were entered into the multiple logistic regression model, and their relationship was assessed in the presence of other variables following the parasitic intestinal infections. The p -values less than 0.05 were considered statistically significant.

Positive/negative parasitic infection was the dependent variable in the logistic regression model. In contrast, other examined variables like sex, age, occupation, education, drinking water source, raw vegetable consumption, and family size were independent or predictive variables.

RESULTS

Our research collected 1038 stool samples from the

Gonabad participants, including 569 (54.8%) males and 469 (45.2%) females. Nearly 31 (3%) of the studied population were illiterate, 180 (17.3%) had primary education, 153 (14.7%) belonged to middle school education level, 452 (43.5%) had completed a diploma, and the rest of the people attained an academic level. Our data showed that 585 (56.4%), 386 (37.2%), and 67 (6.5%) of the studied participants consumed tap water, purified water, and groundwater as water resources, respectively (**Table I**).

Table II demonstrates the prevalence of intestinal parasites detected by the formalin ether sedimentation method. Nearly 18.1% of the total examined stool

samples were positive for at least one species of the parasitic agent out of ten parasitic agents detected in current research.

Our findings indicated that the prevalence of the protozoan parasites (16.7%, 95% CI 14.14 to 19) was more than that of helminthic parasites (1.3%, 95% CI 0.7 to 2). The most prevalent detected parasites were *Blastocystis* sp (12.4%, 95% CI 10.4 to 14.6), *Entamoeba coli* (1.8%, 95% CI 1.1 to 2.6) and *Enterobius vermicularis* (1.2%, 95% CI 0.6 to 1.9). Moreover, as shown in Table.2 *Giardia lamblia* (1.1%), *Entamoeba histolytica /dispar* (0.3%), *Iodamoeba butschlii* (0.3%), *Chilomastix mesnili* (0.3%), *Hymenolepis nana* (0.2 %) *Dientamoeba fragilis*

Table I: Socio-demographic data of participants

Characteristics	Levels	Number (1038)	Percent	Number of infections (187)	Percentage of infection
Sex	Male	569	54.8	87	18.6
	Female	469	45.2	100	17.6
Age	≤ 9	28	2.7	11	39.3
	10 to 19	179	17.2	35	19.6
	20 to 29	334	32.2	54	16.2
	30 to 39	339	32.7	63	18.6
	40 to 49	123	11.8	21	17.1
	≥50	35	3.4	3	8.6
Occupation	Student University	198	19.1	27	13.6
	Student	175	16.9	38	21.7
	Employed	42	4	5	11.9
	Housewife	51	4.9	7	13.7
	Free-Farmer-Livestock	53	5.1	14	26.4
	Labor	466	44.9	89	19.1
	Unemployed	40	3.9	3	7.5
Education	children	13	1.2	4	30.8
	Illiterate	31	3	6	19.4
	Primary	180	17.3	31	17.2
	guidance	153	14.8	31	20.3
	Diploma	452	43.5	81	17.9
	Super Diploma	53	5.1	9	17
	Bachelor	84	8.1	17	20.2
Residency	MA-PHD	85	8.2	12	14.1
	Urban	254	24.5	29	11.4
Source of drinking water	Village	784	75.5	158	20.2
	Tap water	585	56.4	109	18.6
	Purified water	386	37.2	55	14.2
Raw/Unwashed vegetable consumption	Groundwater(qanat)	67	6.5	23	34.3
	No	754	72.6	126	16.7
Family Size	Yes	284	27.4	61	21.5
	Small	350	33.7	62	33.2
	Large	688	66.3	125	66.8

(0.2%) and *Trichomonas hominis* (0.3%) were other observed intestinal parasitic agents. We provided images of trichrome-stained *Entamoeba coli*, *Giardia lamblia* and *Entamoeba histolytica /dispar* in **Figure II**.

Table III exhibited the analyzed data by unadjusted and adjusted logistic regression of the socio-demographic correlates with parasitic intestinal infections among the participants.

Our findings indicated that among possible risk factors investigated in this study, age, drinking water source and residency (rural/urban) status have a significant association with parasitic intestinal infections (p-value < 0.05). In contrast, sex, occupation, education level, raw/unwashed vegetable consumption and the number of family members showed no effects on the rate of parasitic intestinal infections (p-value >0.05)

Table II: Prevalence of intestinal parasites in the studied population

Charac- teristic	Levels	Number	Percent	95% Confidence Interval	
				Lower	Upper
Infection	Blastocystis hominis	129	12.4	10.4	14.6
	Entamoeba coli	19	1.8	1.1	2.6
	Giardia lamblia	11	1.1	0.5	1.8
	Entamoeba. histolytica/ dispar	3	0.3	0.0	0.7
	Dientamoeba fragilis	2	0.2	0.0	0.5
	Trichomonas hominis	3	0.3	0.0	0.7
	Iodamoeba butschlii	3	0.3	0.0	0.7
	chilomastix mesnili	3	0.3	0.0	0.7
	Enterobius vermicularis	12	1.2	0.6	1.9
	Hymenolepis nana	2	0.2	0.0	0.7
	Negative	851	82	80.9	85.4

DISCUSSION

Intestinal parasitic infections are still a significant threat to developing countries. Investigating different localities is crucial to determining high-risk communities and designing appropriate intervention mechanisms. Based on this view, we attempt to assess the prevalence of parasitic intestinal infections and associated risk factors among the human population of Gonabad city, southeast of Khorasan Razavi province, northeastern Iran.

In the present study, the overall prevalence of intestinal parasitic infection was 18.1% among the studied population. These findings more or fewer concur with several previous Iranian studies conducted in different areas like Jiroft city, Khuzestan, Mazandaran, Kohgiluyeh and Boyer-Ahmad, Lorestan and Hamadan provenances with a prevalence of 28%, 25.4% %, 25%, 37.5%, 32.5 % and 35.1% respectively^{3,5,14-16,19,22,27}. We detect that the protozoan infection rate is higher than the helminthic infection. Similar findings were reported from studies conducted in different parts of Iran and the world^{2,5,7,8,21,25}. These variations could be due to population density, climate changes, sanitary facilities, host-parasite interactions, and study methodology⁹. If we review the status of parasitic prevalence in the last four decades, we can see that Iran's overall frequency of parasitic intestinal infections has demonstrated a descending pattern.

Nevertheless, our research and studies mentioned above show that despite progress in medicine and public health, there still needs to be more strategies for controlling these parasitic agents, particularly protozoans, which are major public health issues even nowadays. This problem's importance is reinforced when most previous reports regarding intestinal parasitic prevalence's actual value are generally more than the obtained values. This difference is usually related to non-standard stool sampling (one stool specimen versus three sequential specimens in three alternate days)²². Another probable reason for this difference is the failure to use specific diagnostic methods for intestinal parasites. Similar to recent studies in Iran, in our survey, *Blastocystis sp*, *Entamoeba coli* and *Enterobius vermicularis* were most prevalent among ten types detected parasitic

Figure II: Microscopic picture of detected parasites from Stool samples stained with trichrome staining (magnification 1000x). A. *Giardia lamblia* trophozoite B. *Entamoeba histolytica /dispar* cyst. C. *Entamoeba coli* cyst




Table III: Univariate and multivariate analysis of parasitic intestinal infections and potential risk factors in the participants

Characteristics	Levels	OR ¹	CI ¹ (95%)	p-value	OR ²	CI ² (95%)	p-value
Sex	Female				-		
	Male	1.06	0.77-1.46	0.68		-	
Age	≤ 9				-		
	10 to 19	2.66	1.14-6.18	0.023	2.57	1.06-6.23	0.035
	20 to 29	3.35	1.48-7.56	0.003	2.80	1.19-6.59	0.018
	30 to 39	2.83	1.26-6.34	0.011	2.45	1.05-5.69	0.037
	40 to 49	3.14	1.28-7.66	0.012	2.97	1.17-7.52	0.022
	≥50	6.90	1.69-28.14	0.007	6.12	1.46-25.55	0.13
Occupation	Student University				-		
	Student	0.56	0.33-0.97	0.15			
	Employed	1.16	0.42-3.23	0.76			
	Housewife	0.99	0.40-2.42	0.98			
	Free	0.44	0.21-0.91	0.28		-	
	Labor	0.66	0.41-1.06	0.19			
	Unemployed	1.94	0.56-6.76	0.29			
Education	children	0.35	0.10-1.23	0.16			
	Illiterate				-		
	Primary	1.15	0.43-3.04	0.77			
	guidance	0.94	0.35-2.50	0.90			
	Diploma	1.09	0.43-2.76	0.84			
	Super Diploma	1.17	0.37-3.68	0.78		-	
Residency	Bachelor	0.94	0.33-2.67	0.91			
	MA - PHD	1.46	0.49-4.30	0.49			
Source of drinking water	Village				-		
	Urban	1.98	1.28-2.99	0.002	2.78	1.73-4.46	<0.001
	Tap water				-		
Raw/Unwashed vegetable consumption	purified water	0.43	0.96-1.96	0.045	0.34	0.18-0.63	0.001
	Groundwater (qanat)	1.37	0.25-0.75	0.003	1.53	1.06-2.20	0.021
Family Size	No				-		
	Yes	1.36	0.96-1.91	0.075	1.31	0.92-1.88	0.129
Family Size	Small				-		
	Large	0.97	0.69-1.35	0.85		-	

agents, with a prevalence of 12.4 %, 1.8% and 1.2%, respectively^{5,16,18,19}. The transmission of these parasites through the fecal-oral route indicates poor sanitation practices in Gonabad. Whereas the mean prevalence of *Blastocystis sp* in Iran's total population ranged from 7.5% to 28.4 %, this value in our report was 12.4 % (95% CI 10.4 to 14.6)^{14,19}. Since pathogenicity, zoonotic status, transmission route, and host-parasite adaption pattern of *Blastocystis sp*. have been controversial, further reliable evidence is required to evaluate the associated risk factors related to this infection. Our data demonstrated that 1.8% of parasitic infections belong to *Entamoeba coli* (95% CI 1.1 to 2.6). Recent surveys on apparently healthy Iranian people estimated that up to 18.1 % of cases were infected with *Entamoeba coli*¹⁵, which reflects a

remarkable downregulated prevalence in Iran over the last decade. However, non-pathogenic parasite agents like *Entamoeba coli* do not cause clinical manifestation/infection and are generally transmitted through the faecal-oral route but are regarded as reliable public hygiene indicators. In the present survey, the *Giardia* prevalence was 1.1 % (95% CI 0.5 to 1.8). Other Iranian population-based surveys on *Giardia* frequency rate have stated relatively similar values like 10.2% in 2008 (Kia et al., 2008), 28.2 and 8.9% in 2009^{3,22}, 17.2% in 2011⁵, 2.2% in 2014¹⁵, 17.4 and 7.8% in 2016^{18,27}, and 1.2% in 2017¹⁴. Whereas remarkably high prevalence (91%) reported from Iran highlighted the zoonotic aspect of this protozoan parasite¹⁰. However, a declining tendency in *Giardia* prevalence has been documented

over the last decade in Iran, but it seems relatively high depending on the target population. The decreasing prevalence pattern of human helminth parasites has also been approved in recent decades in Iran.

Nevertheless, some parasites, especially those transmitted through direct fecal-oral routes, like *Enterobius vermicularis* and *Hymenolepis nana*, are still considered prevalent helminthic agents in Iran¹⁷. Here, the prevalence of *Hymenolepis nana* was 0.2 % (95 % CI 0 to 0.7). Although it has been approved that the prevalence of *Hymenolepis nana* in the human population has declined since 1970, this parasitic agent continues to be common in Iran's tribal and rural regions^{18,19,22}. Our data about *Enterobius vermicularis* with a 1.2 % prevalence is similar to recent reports from Tehran and Kerman provinces, where this value was 0.2 % and 0.3 %, respectively^{1,14}. We evaluated several possible risk factors associated with parasitic intestinal infections in the present investigation. In this regard, our analyzed data confirmed that age, drinking water source, and residency (rural/urban) status predict the prevalence of these infectious agents among our studied community.

It was observed that most infected participants belong to age groups ≤ 9 and 10-19, with 39.3 % and 19.6% prevalence rates. It could be assumed that the risk of acquiring infection also increases with aging. People between 40 and 49 and over 50 were approximately three times and nearly seven times, respectively, more likely to be infected than those under nine years old. Similar to our result, an association between intestinal parasitic infections and age has been reported in different parts of Iran¹⁷; this could result from a better understanding and application of hygiene measures. In contrast, this conclusion was not seen in other provenances like Kerman, Tehran and Mazandaran^{1,14,28}. The logistic regression test demonstrated no association between occupation status and intestinal parasitic infection (p -value >0.05). Most infected people were Free-Farmer-Livestock (26.4%). Similar findings were documented from different parts of Iran, such as the Kerman, Tehran, Mazandaran and Isfahan provinces^{1,14,16,18,19}.

We found no significant association between educational background and infection with intestinal parasites (p -value >0.05). This subject differs from another study from Kurdistan province that showed literacy level is a protective factor against infection⁴. This variation in educated people could be due to variations in awareness about the mode of parasitic agent transmission and applying the necessary measures to avoid it. In contrast, we did not see this pattern in our examination. The most identified parasites belong to the group with a diploma educational level, which could result from a participant ratio at this academic level. Regarding residence, people who came from rural areas had a higher parasitic infection rate (20.2 %) and were approximately three times more at risk (OR = 2.78 ,

%95CI 1.73-4.46 , $p <0.001$) of getting infected with parasitic agents than urban dwellers. Several similar findings have been approved in different parts of Iran. It was proposed that limited access to adequate sanitation, inadequate healthcare services coverage, low socioeconomic status, and hygienic living conditions are the most probable factors that increase the risk of infection in rural people^{4,5,30}. The kind of source of drinking water is a leading factor in the prevalence, incidence and dissemination of pathogen agents, particularly waterborne infections. Interestingly, we found that consuming purified water protects against infection, while underground water consumption increases the risk of infection. In recent years, Gonabad has experienced severe drought and faces significant problems in water resources due to several causes, such as population burden, industrial activities, climate changes, and increasing demand²⁹. Nearly all drinking water resources in arid regions like Gonabad City depend on groundwater supplies⁶ (Bakarman et al., 2019). Additionally, based on traditional belief, underground water resources like qanat have therapeutic benefits. Although the different aspects of Gonabad underground water (like heavy metal content) have been investigated several times, their safety is controversial. Despite that, these water supplies do not seem suitable for drinking water. Comparable results have been regarding the importance of drinking water sources^{14,18}. Previous studies imply that contamination of vegetables with parasitic agents could be the primary cause of infection spread in different parts of Iran. However, our analyzed data does not support this assertion¹⁷. As a unique characteristic of this research, we found no association between family size and parasitic intestinal infection, unlike two other studies in Ethiopia. This difference may be due to sample size and residency status³².

Limitations

There were several limitations in the present study. The most remarkable one is that we could not apply swabs (cellophane-tape) tests to comment on the actual prevalence of *Enterobius vermicularis* due to the large sample size and required condition. Moreover, we cannot apply the molecular approach to detect parasitic agents due to the monetary burden. Moreover, only one stool specimen was used to identify parasites; this approach may lead to underestimating the actual prevalence of intestinal parasitic agents compared to three sequential specimens collected in three alternate days.

CONCLUSION

The present report provides a snapshot of the status of intestinal parasitic prevalence and related factors in Gonabad. Although there has been a decreasing trend in prevalence in Iran, the parasite burden rate of 18.1% in Gonabad is concerning. The report identifies age, residency status, source of drinking water and

raw/unwashed vegetable consumption as potential risk factors for parasitic intestinal infections. The study's findings could help inform policies to improve public health through lifestyle changes, promoting public awareness, and infection prevention and control. The report also recommends periodic assessment and treatment of human cases of parasitic infections.

Acknowledgements: We want to acknowledge all staff from the Vector-Borne Disease Research Center, North Khorasan University of Medical Sciences, Bojnurd, Iran, for their helpful assistance.

Ethical Considerations: Gonabad University of Medical Sciences, Iran ERC Letter No. A-10-172-51). IR.GMU.REC.1398.010.

Financial Disclosure / Grant Approval: The author (s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article. A grant from Gonabad University of Medical Sciences financially supported this study.

Conflicts of Interest: The authors declare no conflict of interest.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

Salehi M: Methodology creation and writing
Razavi S: Manuscript writing, review and editing
Khajavian N: Statistical analysis
Ghasemzadeh-Moghaddam H: Original draft manuscript preparation and writing
Firouzeh N: Manuscript writing, editing and review
Minooeian Haghighi MH: Funding acquisition and writing

REFERENCES

1. Afshar MJ, Barkhori Mehni MB, Rezaeian M, Mohebali M, Baigi V, Amiri S et al. Prevalence and associated risk factors of human intestinal parasitic infections: A population-based study in the southeast of Kerman province, southeastern Iran. *BMC Infect Dis.* 2020; 20(1): 12. doi: 10.1186/s12879-019-4730-8.
2. Arani AS, Alaghebandan R, Akhlaghi L, Shahi M, Lari AR. Prevalence of intestinal parasites in a population in south of Tehran, Iran. *Rev Inst Med Trop Sao Paulo.* 2008; 50(3): 145-9. doi: 10.1590/S0036-46652008000300003.
3. Badparva E, Kheirandish F, Ebrahimzade F. Prevalence of intestinal parasites in Lorestan Province, West of Iran. *Asian Pacific J Trop Dis.* 2014; 4(2): S728-S732. doi: 10.1016/S2222-1808(14)60716-7.
4. Bahrami F, Haghighi A, Zamini G, Khadem-Erfan MB, Azargashb, E. Prevalence and associated risk factors of intestinal parasitic infections in Kurdistan province, Northwest Iran. *Cogent Med.* 2018; 5(1): 1503777. doi: 10.1080/2331205x.2018.1503777.
5. Kuzehkanani AB, Rezaei S, Babaei Z, Niyati M, Hashemi SN, Rezaeian M. Enteric protozoan parasites in rural areas of Bandar-Abbas, southern Iran: Comparison of past and present situation. *Iran J Public Health.* 2011; 40(1): 80-85.
6. Bakarman MA, Hegazi MA, Butt NS. Prevalence, characteristics, risk factors, and impact of intestinal parasitic infections on school children in Jeddah, western Saudi Arabia. *J Epidemiol Glob Health.* 2019; 9(1): 81-87. doi: 10.2991/jeqh.k.190219.001.
7. Boonjaraspinyo S, Boonmars T, Kaewsamut B, Ekobol N, Laummaunwai P, Aukkanimart R et al. A cross-sectional study on intestinal parasitic infections in rural communities, northeast Thailand. *Korean J Parasitol.* 2013; 51(6): 727-34. doi: 10.3347/kjp.2013.51.6.727.
8. Calderaro A, Montecchini S, Rossi S, Gorrini C, De Conto F, Medici MC et al. Intestinal parasitoses in a tertiary-care hospital located in a non-endemic setting during 2006-2010. *BMC Infect Dis.* 2014; 14: 264. doi: 10.1186/1471-2334-14-264.
9. Elmonir W, Elaadli H, Amer A, El-Sharkawy H, Bessat M, Mahmoud SF et al. Prevalence of intestinal parasitic infections and their associated risk factors among preschool and school children in Egypt. *PLoS ONE.* 2021; 16(9): e0258037. doi: 10.1371/journal.pone.0258037.
10. Faridi A, Kareshk AT, Sadooghian S, Firouzeh N. Frequency of different genotypes of *Giardia duodenalis* in slaughtered sheep and goat in east of Iran. *J Parasit Dis.* 2020; 44(3): 618-624. doi: 10.1007/s12639-020-01237-1.
11. Fauziah N, Aviani JK, Agrianfanny YN, Fatimah SN. Intestinal parasitic infection and nutritional status in children under five years old: a systematic review. *Trop Med Infect Dis.* 2022; 7(11): 371. doi: 10.3390/tropicalmed7110371.
12. Girma A, Aemiro A. Prevalence and Associated Risk Factors of Intestinal Parasites and Enteric Bacterial Infections among Selected Region Food Handlers of Ethiopia during 2014-2022: A Systematic Review and Meta-Analysis. *Scientif World J.* 2022; 2022: 7786036. doi: 10.1155/2022/7786036.
13. Hajissa K, Islam MA, Sanyang AM, Mohamed Z. Prevalence of intestinal protozoan parasites among school children in Africa: A systematic review and meta-analysis. *PLoS Negl Trop Dis.* 2022; 16(2): e0009971. doi: 10.1371/journal.pntd.0009971.
14. Hemmati N, Razmjou E, Hashemi-Hafshejani S, Motevalian A, Akhlaghi L, Meamar AR. Prevalence and risk factors of human intestinal parasites in Roudehen, Tehran province, Iran. *Iran J Parasitol.* 2017; 12(3): 364-373.
15. Jafari R, Fallah M, Darani HY, Yousefi HA,

- Mohaghegh MA, Latifi M et al. Prevalence of Intestinal Parasitic Infections Among Rural Inhabitants of Hamadan City, Iran, 2012. *Avicenna J Clin Microbiol Infect.* 2014; 1(2): 21445. doi: 10.17795/ajcmi-21445.
16. Kia EB, Hosseini M, Nilforoushan MR, Meamar AR, Rezaeian M. Study of intestinal protozoan parasites in rural inhabitants of Mazandaran Province, northern Iran. *Iran J Parasitol.* 2008; 3 (1): 21-25.
 17. Mahmoudvand H, Badparva E, Khalaf AK, Niazi M, Khatami M, Nazer MR. Prevalence and associated risk factors of intestinal helminthic infections in children from Lorestan province, Western Iran. *Parasite Epidemiol Control.* 2020; 9: e00136. doi: 10.1016/j.parepi.2020. e00136.
 18. Mahni MB, Rezaeian M, Kia EB, Raeisi A, Khanaliha K, Tarighi F et al. Prevalence of intestinal parasitic infections in Jiroft, Kerman Province, Iran. *Iran J Parasitol.* 2016; 11(2): 232-238.
 19. Mowlavi GR, Mirahmadi H, Rezaeian M, Kia E, EbrahimiDaryani N, Rokni MB et al. Prevalence of intestinal parasites in tribal parts of Khuzestan Province during 2005-07. *Govaresh.* 2008; 12(4): 219-228.
 20. Ngwese MM, Manouana GP, Moure PAN, Ramharter M, Esen M, Adégnika AA. Diagnostic techniques of soil-transmitted helminths: Impact on control measures. *Trop Med Infect Dis.* 2020; 5 (2): 93. doi: 10.3390/tropicalmed5020093.
 21. Niyiyati M, Rezaeian M, Zahabion F, Hajarzadeh R, Kia EB. A survey on intestinal parasitic infections in patients referred to a hospital in Tehran. *Pak J Med Sci.* 2009; 25(1): 87-90.
 22. Pestehchian N, Nazari M, Haghighi A, Salehi M, Yosefi H, Khosravi N. Prevalence of intestinal parasitic infection among inhabitants and tribes of Chelgerd, Iran, 2008-2009. *J Clin Diagn Res.* 2015; 9(5): LC01-LC04. doi: 10.7860/JCDR/2015/11580. 5879.
 23. Qasemi M, Afsharnia M, Farhang M, Bakhshizadeh A, Allahdadi M, Zarei A. Health risk assessment of nitrate exposure in groundwater of rural areas of Gonabad and Bajestan, Iran. *Environ Earth Sci.* 2018; 77: 551. doi: 10.1007/s12665-018-7732-8.
 24. Rasti S, Hassanzadeh M, Hooshyar H, Momen-
Heravi M, Mousavi SGA, Abdoli A. Intestinal parasitic infections in different groups of immunocompromised patients in Kashan and Qom cities, central Iran. *Scand J Gastroenterol.* 2017; 52(6-7): 738-741. doi: 10.1080/00365521.2017.1308547.
 25. Rokni MB. The present status of human helminthic diseases in Iran. *Ann Trop Med Parasitol.* 2008; 102(4): 283-95. doi: 10.1179/136485908X300805.
 26. Saki J, Amraee D. Prevalence of Intestinal Parasites Among the Rural Primary School Students in the West of Ahvaz County, Iran, 2015. *Jentashapir J Health Res.* 2017; 8(1): e40326. Doi: 10.5812/jjhr.40326.
 27. Sarkari B, Hosseini G, Motazedian MH, Fararouei M, Moshfe A. Prevalence and risk factors of intestinal protozoan infections: A population-based study in rural areas of Boyer-Ahmad district, Southwestern Iran. *BMC Infect Dis.* 2016; 16(1): 703. doi: 10.1186/s12879-016-2047-4.
 28. Shahdoust S, Niyiyati M, Haghighi A, Azargashb E, Khataminejad MR. Prevalence of intestinal parasites in referred individuals to the medical centers of Tonekabon city, Mazandaran province. *Gastroenterol Hepatol Bed Bench.* 2016; 9 (Suppl1): S75-S79.
 29. Sitotaw B, Shiferaw W. Prevalence of Intestinal Parasitic Infections and Associated Risk Factors among the First-Cycle Primary Schoolchildren in Sasiga District, Southwest Ethiopia. *J Parasitol Res.* 2020; 2020: 8681247. doi: 10.1155/2020/8681247.
 30. Symeonidou I, Tassis P, Gelasakis AI, Tzika ED, Papadopoulos E. Prevalence and risk factors of intestinal parasite infections in Greek swine farrow-to-finish farms. *Pathogens.* 2020; 9(7): 556. doi: 10.3390/pathogens9070556.
 31. Taherkhani K, Barikani A, Shahnazi M, Saraei M. Prevalence of intestinal parasites among rural residents of Takestan in north-west of Iran. *Iran J Parasitol.* 2019; 14(4): 657-663. doi: 10.18502/ijpa.v14i4.2118.
 32. Tsegaye B, Yoseph A, Beyene H. Prevalence and factors associated with intestinal parasites among children of age 6 to 59 months in, Boricha district, South Ethiopia, in 2018. *BMC Pediatr.* 2020; 20 (1): 28. doi: 10.1186/s12887-020-1935-3.

