# Awareness of Dental Undergraduates, Post Graduates and Dental Practitioners' about Dental and Biomedical Waste Management

Muhammad Abdullah Kamran, Uzma Zareef, Tauseef Ahmed, Raffat Rasool, Nabeel Khan, Mehwash Kashif

# ABSTRACT

OBJECTIVE: To assess undergraduate, postgraduate dental students' and dental practitioners' knowledge of management of dental and biomedical waste and to focus on critical issues such as environment-friendly waste management.

METHODOLOGY: A total of 273 participants were recruited from January to June 2019 through convenience sampling in this cross-sectional study; undergraduate, postgraduate dental students, and general dental practitioners from private and public sector dental colleges of Karachi. The respondents answered different categories of waste management, including biomedical waste rules, waste disposal measures, dental waste types, common dangers of improper waste management, and some particular equipment problems, data analyzed on SPSS version 17.00.

RESULTS: From 273 (92.8%) completed questionnaires, 217(79.4%) were females. Undergraduate students were 61.9%, (n=169). Almost 64.4 % (n=176) of respondents didn't know about dental waste management rules. Among 75 % (n=205) of respondents, there was a consensus that qualified professionals should treat hospital disposal. Only 67.7% (n= 185) of individuals surveyed stated that they were familiar with the categories formed by dental waste. Approximately 95.5% (n=260) of respondents recommend workshops and Continues medical education seminars for waste management. Compared to undergraduate students, postgraduate students had a significantly higher mean score, decreasing the "know-do gap" (p=0.069).

CONCLUSION: Among undergraduate dental students and dental practitioners, postgraduates were found to have a greater understanding of the correct methods for disposing of healthcare waste. However, the overall knowledge level of environment friendly waste management was insufficient.

## KEYWORDS: Disposal of medical waste; Understanding of health, attitudes, practice; Dentistry

*This article may be cited as:* Kamran MA, Zareef U, Ahmed T, Rasool R, Khan N, Kashif M. Awareness of Dental Undergraduates, Post Graduates and Dental Practitioners' about Dental and Biomedical Waste Management. J Liaquat Uni Med Health Sci. 2022;21 (01):50-4. doi: 10.22442/jlumhs.2022.00864. Epub 2022 January 12.

## INTRODUCTION

The health care system is a major profession in Pakistan. It includes various health sectors like medicine, surgery, dentistry, pharmacy, physiotherapy, etc. Waste generated by these health care domains is insufficient quantity<sup>1</sup>. Healthcare waste encompasses all of the garbage that healthcare services, testing services, and research labs generate<sup>1</sup>. Moreover, waste from dispersed resources generated in-home health care, like dialysis supplies, insulin injections, etc., was too incorporated<sup>2</sup>. Approximately more than 80% of hospital waste is not harmful, among which WHO estimates; about 10% is contagious, and the remaining 5% is non-infectious. Awareness regarding the proper management of medical and dental waste products, especially infectious ones, is of utmost importance for the appropriate care of patients, people, and the environment<sup>3</sup>.

Most dental professionals seek to function independently inside their clinics; they must be aware of proper waste management. To fulfill this requirement, it must be taught as part of the dental curriculum during undergraduate clinical training<sup>4</sup>, like hepatitis and HIV infections, improper waste treatment results in the aggressive spread of infection. Perhaps it serves as a breeding place for numerous other disease transmission vectors<sup>5</sup>. The spread of such infections from waste from health care presents a danger to people's health and their climate. It is necessary to emphasize the proper management of biomedical waste (BWM)<sup>6</sup>. Only a few studies have been conducted in the last five years regarding waste management in dental practices in Pakistan<sup>7</sup>. Several of these researches highlighted the immediate demand for new education in BWM technologies and suggested proper disposal of biomaterials and radiography waste which may have a more negligible effect on the surroundings. Research on biomedical management awareness and attitudes has also highlighted insufficient knowledge. substantial variance, and the call for education courses<sup>8</sup>.

As a result, this study designed to assess undergraduate, postgraduate dental students and

### Dental Practitioners' about Dental and Biomedical Waste Management

dental practitioners' knowledge regarding dental and biomedical waste management and focus on critical issues such as environment-friendly waste management.

# METHODOLOGY

Cross-sectional research was planned among 273 respondents from January to June 2019 through nonprobability convenience sampling; subjects were recruited for the study. The study conducted among undergraduates, postgraduate students and general dental practitioners of Karachi Medical and Dental College and Liaguat College of Medicine and Dentistry, Karachi. The sample size was calculated using Raosoft software with the accepted margin of error of 5%, 95% confidence level, the population size of 930, and response distribution of 50%. After taking informed verbal consent, participants were asked to fill out a closed-ended questionnaire about the researchers' knowledge of BWM processes. There were five domains in the questionnaire, each of which evaluated different aspects of BWM in dentistry. A research questionnaire was comprised of 25 questions from five areas, such as "ethical aspects of biomedical waste", "stages of waste management", "categories of dental care waste", "common hazards of inappropriate waste disposal," and specific relevant equipment questions" was used to collect data. They were asked to choose the most suitable response and were told that their anonymity was confidential.

SPSS program 17.00 (SPSS Inc., III., USA) was used to conduct statistical analyses. The overall score of the questionnaire was calculated by giving accurate or acceptable responses a value of '1' and a score of '0'. The overall score was compared between the sexes, designations, and qualifications of the participants using the Mann-Whitney U test, which compares the overall score among sexes, designations, and qualifications of the participants. At 0.05, the p-value was determined to be statistically significant.

# RESULTS

Approximately 273 (92.8%) complete questionnaires were chosen for data analysis. Total 217 (79.4%) were female among the participants. Most of the participants were undergraduate students (61.9%, n=169), while 17.3% (n=49) were postgraduates, and 20.14% (n=55) were general practitioners. Overall, 64.4 % (n=176) of respondents didn't know about dental waste management rules in Pakistan (Table I). Approximately 53.1% (n=145) participants responded that waste could be kept up to 96 hours before disposal, while 19.4% (n=53) reported 12 hours to dispose of the waste. Just 9.15 % (n=25) suggested that 48 hours was the highest allowable period for the waste stores. Most respondents (67.0%, n=183) noted that a government-approved collector is responsible for the final disposal of dental wastage. (Table I)

### TABLE I: SHOWING AWARENESS OF BIOMEDICAL WASTE MANAGEMENT IN PAKISTAN

Query	Response	n	%
Awareness of Pakistan's	Yes	31	11.3
biomedical waste	No	176	64.4
management laws?	Not certain	66	24.1
Duration of waste keeping	12 hr	53	19.4
	72 hr	25	9.15
	48 hr	145	53.1
	96 hr	50	0.0
Awareness of the dental	Yes	18	6.59
waste transportation	No	72	26.3
regulatory body	Don't know	183	67.0

The second area (**Table II**) consisted of knowledge about stages of waste management. Approximately (53.8 %, n= 147) perceived that they were sufficiently conscious of the skills needed to manage hospital waste. Among 75 % (n=205) of respondents, it recommends that qualified professionals should treat hospital disposal. Results regarding awareness regarding proper order of six successful waste management steps revealed that only 42.8% (n= 117) respondents know the appropriate steps of waste disposal, i.e., Collection- Segregation- Storage-Transportation- Disposal.

The different types of waste generated in various dental specialty departments were assessed in the third category (**Table III**); this covered approximately one-third of the questions due to its special significance and relevance in everyday practice. About 67.7% (n= 185) of individuals surveyed stated they were not familiar with the categories formed by dental waste. The majority of respondents, i.e., 48.3% (n=132), consider extracted teeth can be infected or non-infected. Pharmaceutical waste is deemed the significant component of waste management, i.e., 64.1 % (n= 175). The majority of respondents (60.4. n= 129) consider a group of second-hand impression materials and cotton as highly contaminated items. Amalgam can be stocked up in common trash was responded by 53.1(n=153) respondents.

53.1 % (n= 145) of respondents can dispose of used sharps in puncture-resistant plastic bags. Discarding developer and fixer solution is majorly poured down the drain after being diluted, as responded by 65.5 (n=179) respondents. About 58.9% (161) respondents don't know about the disposal of exposed X-ray Films. The final group (**Table IV**) consisted of five general information-based questions about the various hazards and the value of waste disposal, which are typically taught in the BDS curriculum. Many of the questions in this section were correctly answered. Still, most respondents (95.2%, n=260) recommend

Muhammad Abdullah Kamran, Uzma Zareef, Tauseef Ahmed, Raffat Rasool, Nabeel Khan, Mehwash Kashif

# TABLE II: KNOWLEDGE OF BIOMEDICAL WASTE MANAGEMENT

Query	Reaction   Yes   No   Not sure		% 25.2 53.8 20.8
Awareness regarding the handling of hospital waste?			
Biomedical waste handling by trained workers	No Yes Can't Say	10 205 58	3.6 75 21.2
Awareness regarding proper order of six successful waste management steps?	Segregation-Collection- Transportation- Storage- Treatment- Disposal Collection- Segregation- Storage- Transportation- Disposal	89 117	30.03 42.8
•	Storage- Segregation- Transportation- Collection- Treatment - Disposal	45	16.4
•	Transportation- Segregation- Collection- Storage- Treatment - Disposal	22	8.0

### TABLE III: QUESTIONS RELATED TO AWARENESS OF WASTE MANAGEMENT

Factors	Reaction	n	%
Awareness of dif- ferent categories of biomedical wastes	Don't know Know	185 88	67.7 32.2
Extracted teeth can be categorized into	Infected Non-infected Both Don't know	100 21 132 20	42.1 7.6 48.3 7.3
Disposal of a group of obsolete drugs	Chemical waste Pharmaceutical waste Cytotoxic waste Don't know	56 175 27 15	20.5 64.1 9.8 5.4
Group of second- hand impression materials and cotton	Cytotoxic dirty contaminated Don't distinguish	10 98 129 36	3.6 35.8 60.2 13.1
Amalgam can be stocked up	Place in a common trash can. Only airtight containers are al- lowed. Water in an airtight container Using the fixer	115 67 49 42	53.1 15.3 17.9 55.9
Disposal of used sharps	A typical bin Puncture-resistant plastic bag Break the needle and toss it away Incineration	26 145 95 7	9.5 53.1 34.7 2.5
Discarding of de- veloper and fixer solution	Both of them end up in the sewer. The silver from the fixer is recov- ered and returned to the source in a designated facility	8 49	2.9 17.9
	Poured down the drain after being diluted Others	179 45	65.5 16.4
Disposal of ex-	Stored separately and	96	35.1
posed X-ray Films	disposed Buried in soil Don't know	16 161	5.8 58.9

waste management workshops and teaching so that realistic waste management exercises in dentistry schools would be beneficial. The questionnaire's overall score was determined by assigning a score of '1' for accurate or approved responses and a score of '0' for erroneous replies. Table V reveals that the mean total score values for designation, and qualification were sex, not significantly different (p=0.413 and p=0.076, respectively). As compared to students, postgraduate students had a significantly higher mean score, decreasing the "know-do gap" (p=0.069). The response given was that postgraduates spend more time in the hospital.

## TABLE IV: THE DANGERS OF INCORRECT

Query	Answer	n	%
Regarding	Should be kept closed	119	43.5
biomedical waste	Should be clean in the open air	45	16.4
containers	Compatibility according to waste	32	11.7
	Type of container used	77	28.2
Contribution to pollution	Foul odor and growth of insects	28	10.2
and disease Spread of Disease		127	46.5
dispersal	Both	109	39.9
Disinfection	Yes	178	65
of Lead	No	29	6.5
Aprons	Don't know	66	30.1
Hazardous	Silver	15	5.4
	Tin and Copper	5	1.8
amalgam?	Zinc	11	4.0
	Mercury	242	88.6
Recommen-	Agree	260	95.2
dations for waste man- agement workshops and teaching	Disagree	13	4.7

PARTICIPANTS					
Factors	Category —		Score		
racions		n	Mean	± SD	value
Gender	Women	217	11.70	2.96	0.413
	Men	56	13.96	2.35	0.415
Designation	Undergraduate	169	12.46	2.83	
	Postgraduate	64	15.22	2.74	0.069
	GDPs	40	13.8	2.89	
Qualification	BDS	55	10.58	2.81	
	FCPS/MDS	49	11.17	2.80	0.076
	Undergraduate	169	11.59	3.50	

# TABLE V: DEMOGRAPHIC DETAILS OF STUDY

# DISCUSSION

The results show a few gaps in undergraduate dental students, postgraduate students, and general dental practitioners regarding dental healthcare waste management regulations and awareness<sup>9,10</sup>. However, a limited number of waste management experiments have been conducted at dental schools<sup>11,12,16</sup>.

Those aspects are unfamiliar to dental students and students. They are not taught as part of the curriculum, so students have no idea what happens to the dental materials they use or the many options for adequately recycling or reusing them. Sharps are thrown away in a standard container after breaking the needle, which is a big concern and inappropriate way to manage sharps, as indicated by around 53.1% of participants in this study's comments; this is similar to other research in which 40% and 58 percent of respondents obtained the same answers, indicating a significant "know-do gap"<sup>17,18</sup>.

It is critical to establish fines/penalties for the incorrect disposal of dental healthcare waste on a regulatory level and teach dental undergraduates about this in their early years of dentistry<sup>20</sup>.

This research had a few limitations because only dentistry students were recruited, rather than the auxiliary personnel who is the driving force behind the BWM. Program limitations may have been reflected as a disparity in the participants' awareness of the BWM; dental auxiliary should also be assessed for the knowledge and handling of dental waste. Crosssectional design and small sample s size were also the study's limitations. It is recommended that further studies be planned with large sample size and modify study design to generalize results.

The main reason for this is a lack of education, which only addresses the disposal of biomedical waste in the clinical setting and offers little knowledge on the negative environmental consequences of improper disposal and modern proper disposal technologies. In comparison to the course, studies have shown that dental practitioners and students would participate in continuing dental education. It resulted that everyone working in the healthcare sector has a responsibility to reduce the risks in society. This can be accomplished by educating students as early as at the undergraduate level, informing them of the most recent advancements in waste disposal, and including of efficient the practical application waste management into the dental curriculum. It is necessary to improve global awareness of proper waste disposal methods and introduce modern waste recycling methods where possible. The areas of our curriculum that need to be improved have been highlighted in this article. The need for changes and upgrades has been expressed so that future generations of dentists will minimize the burden of improper Waste management practices in health care with our environment and disease transmission.

It is advised that waste management teaching should be included in the dental curriculum. Dental personnel must also be taught and trained regarding proper and appropriate disposal of biomedical waste; workshops and courses should be planned to educate people to prevent harmful effects of waste management. It is also recommended to draw attention to educational syllabus flaws and curriculum, emphasizing ecofriendly waste management.

# CONCLUSION

Among undergraduate dental students and dental practitioners, postgraduates were found to have a greater understanding of the correct methods for disposing of healthcare waste. However, the overall knowledge level of environment-friendly waste management was insufficient.

**Ethical permission**: Liaquat College of Medicine & Dentistry, Karachi, letter Dated: 24-12-2018.

**Conflict of Interest:** There is no conflict of interest among the authors.

**Financial Disclosure / Grant Approval:** There was no funding agency.

**Data Sharing Statement**: The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions

## AUTHOR CONTRIBUTIONS

Kamran MA: Concept, data collection, write-up review Zareef U: Concept, write-up, data collection Ahmed T: Data collection, critical review, analysis Rasool R: Write-up, data collection, analysis Khan N: Write-up, analysis, data collection Kashif M: Write-up, analysis, critical review

# REFERENCES

 Kumar R, Khan EA, Ahmed J, Khan Z, Magan M, Nousheen N et al. Healthcare waste management (HCWM) in Pakistan: current situation and training options. J Ayub Med Coll Abbottabad. 2010; 22 (4): 101-5.

### Muhammad Abdullah Kamran, Uzma Zareef, Tauseef Ahmed, Raffat Rasool, Nabeel Khan, Mehwash Kashif

- Ishtiaq P, Khan SA, Haq MU. A multi-criteria decision-making approach to rank supplier selection criteria for hospital waste management: A case from Pakistan. Waste Manag Res. 2018; 36(4): 386-94.
- 3. Organization WH. Management of waste from hospitals and other health care establishments: Report on a WHO meeting. EURO Rep Stud. 1985; 97: 1-61.
- Tenna A, Stenehjem EA, Margoles L, Kacha E, Blumberg HM et al. Infection control knowledge, attitudes, and practices among healthcare workers in Addis Ababa, Ethiopia. Infect Control Hosp Epidemiol. 2013; 34(12): 1289-96. doi: 10.1086/673979.
- Organization WH. Preparation of National Health-Care Waste Management Plans in Sub Saharan Countries. Guidance Manual. Available from: https://apps.who.int/iris/bitstream/handle/ 10665/43118/924154662X.pdf
- Puri S, Smriti K, Pentapati KC, Singh R, Vineetha R, Tamrakar A. Assessment of Awareness About Various Dental Waste Management Practices Among Dental Students and Practicing Clinicians. Pesquisa Brasileira em Odontopediatria e Clínica Integrada. 2019; 19(1): e4839. doi: 10.4034/ PBOCI.2019.191.136.
- Kumar R, Somrongthong R, Ahmed J. Effect of medical waste management trainings on behavior change among doctors versus Nurses and paramedical staff in Pakistan. J Ayub Med Coll Abbottabad. 2016; 28(3): 493-6.
- Kapoor D, Nirola A, Kapoor V, Gambhir RS. Knowledge and awareness regarding biomedical waste management in dental teaching institutions in India - A systematic review. J Clin Exp Dent. 2014; 6(4): e419-e424. doi: 10.4317/jced.51565.
- Abhishek KN, Supreetha S, Varma Penumatsa N, Sam G, Khanapure SC, Sivarajan S. Awarenessknowledge and practices of dental waste management among private practitioners. Kathmandu Univ Med J. 2016; 14(53): 17-21.
- Organization WH. United Nations Environment Programme / SBC. National Health-Care Waste Management Plan - Guidance Manual. Fundamentals of healthcare waste management. Available from: https://www.who.int/water\_ sanitation\_health/medicalwaste/en/guidance manual1.pdf.

## AUTHOR AFFILIATION:

**Dr. Muhammad Abdullah Kamran** Professor, Department of Orthodontics Liaquat College of Medicine and Dentistry Dar-ul-Sehat Hospital (LCMD, DSH) Karachi, Sindh-Pakistan.

## Dr. Uzma Zareef

Professor, Department of Oral Pathology LCMD, DSH, Karachi, Sindh-Pakistan.

## Dr. Tauseef Ahmed,

Assistant Professor Department of Oral Pathology LCMD, DSH, Karachi, Sindh-Pakistan.

## Dr. Raffat Rasool

Assistant Professor Department of Forensic Medicine Karachi Medical and Dental College Karachi, Sindh-Pakistan.

- Kahar AR, Arora A, Radke U, Joshi J. Assessment of awareness regarding biomedical waste management among students and interns of dental institute. Indian J Multidiscip Dent 2017; 7(2): 65-70.
- 13. American Dental Association. Best management practices for amalgam waste. J Okla Dent Assoc. Summer 2004; 95(1): 28-30.
- 14. Ranjan R, Pathak R, Singh DK, Jalaluddin M, Kore SA, Kore AR. Awareness about biomedical waste management and knowledge of effective recycling of dental materials among dental students. J Int Soc Prev Community Dent. 2016; 6 (5): 474-9.
- Oregon Dental Association. The Environmentally Responsible Dental Office: A Guide to Best Management Practices of Dental Wastes. 2008; 1 -12. Available from: https://www.eugene-or.gov/ DocumentCenter/View/4978/Best-Management-Practices--Dental-Offices
- Arora S, Mittal S, Dogra V. Eco-friendly dentistry: Need of future. An overview. J Dent Allied Sci. 2017; 6: 22-7.
- 17. NHS UK. (HTM 07-01) Management of disposal of healthcare waste. 2013. Available from: https:// www.england.nhs.uk/publication/management-and-disposal-of-healthcare-waste-htm-07-01/.
- Prouse RE, West AA, King DA, Poulson R. Method of making paper from water-insoluble alginate fibers and the paper produced. Available from: https://patentimages.storage.googleapis. com/f6/62/5b/0fc32fd621bca1/US4104115.pdf.
- Lakshmikantha R, Kanyadara J, Bullappa D, Vanishree N, Prasad KSK, Naveen N et al. To assess the knowledge, level of awareness, and attitude on biomedical waste management among practicing dentists in Bengaluru city: A crosssectional study. CHRISMED J Heal Res 2016; 3 (3):161-7.
- Ladia S, Gupta N. Effectiveness of training program related to infection control and waste management practices in a private dental college, Pune - A quasi-experimental study. J Indian Assoc Public Heal Dent 2017; 15(1): 4-7.

#### Dr. Nabeel Khan Assistant Professor Department of Oral Biology Karachi Medical and Dental College Karachi, Sindh-Pakistan. Dr. Mehwash Kashif (Corresponding Author) Professor, Department of Oral Pathology Karachi Medical and Dental College Karachi, Sindh-Pakistan. Email: mehwashkashif@gmail.com