

Health Consumer Expectations and Perception of Quality Care Services at Primary Health Care Level in Nigeria

Saka Mohammed Jimoh, Akande Tanimola Mankanjuola, Saka Aishat Oluwatoyin, Bello Fatai. A, Isiaka Sulu Babaita

ABSTRACT

OBJECTIVE: The study examined the expectations of health consumers and perception used in judging the Quality of health care services at the primary health care level.

METHODOLOGY: A cross-sectional study was carried out among randomly selected health consumers that had received health care services in the hospitals designated as primary health care facilities in Kwara State Nigeria.

RESULTS: Overall health consumers' means score expectations was 6.57 while perception was 5.80. Though was high in favour of females with no statistics significant difference. However, correlation test revealed significant association in socio-demographic variables such as age, occupation and educational status. Conversely, relationship was inverse in term of increase in education attainment with lower perception value ($pV = <0.001$).

CONCLUSIONS: Expectation- perception gap was a major determinant of quality of health care services at primary health care levels. Significant relationship occurred between health consumer's age, occupation, educational status and overall expectation -perception of quality health care received. It is recommended that hospital management need to monitor the link between the expectations including perception of services received by patients for quality improvement at primary health care level.

KEY WORDS: Health Consumer Expectations, Quality Care Services, Perception, Nigeria

This article may be cited as: Jimoh SM, Mankanjuola AT, Oluwatoyin SA, Fatai BA, Babaita IS. Health Consumer Expectations and Perception of Quality Care Services at Primary Health Care Level in Nigeria. J Liaquat Uni Med Health Sci. 2019;18(01):47-54. doi: 10.22442/jlumhs.191810600

INTRODUCTION

Quality of health care involves offering technical competent, effective and safe care that contributes to the client's well-being, it is doing the right thing right, right away so as to meet the health consumer expectations, obtaining the best possible clinical outcome and maintaining positive perception of services¹.

Expectations, with reference to healthcare, refer to the anticipation or the belief about what is to be encountered in a consultation or in the healthcare system. It is the mental picture that health consumers or the public will have of the process of interaction with the system. Every patient who comes for consultation has expectations based on his understanding of the illness, cultural background, health beliefs, and attitudes, level of understanding and socio-demographic characteristic of the health consumers. How far the physician reaches an understanding with the health consumers will also have an impact on the successful outcome of the consultation². Not meeting expectations can result in non-compliance, the health consumers may never complain to the physician directly but instead they just

will not return for on-going and follow-up care^{3,4}. Perceptions mean how health plans, hospitals, physicians, and health care in general are viewed currently by the health consumers. It is affected by (a) the dominance of market-oriented approaches to reforms in health care delivery and cost; and (b) the emergence of a normative perspective on clinical practice that emphasizes the need to deliver health consumer-centered care⁵⁻¹⁰. There are two perspectives on the important of patient perceptions of quality of health care, firstly, one can be normative that patient perceptions of quality are inherently meaningful and should be a primary focus of attention within the health care system. Secondly, one can take the position that we have to pay attention to patient perceptions because they are powerful drivers of outcomes important to various other stakeholders, outcomes such as patient choice of plan or provider, patient adherence to medical advice,¹¹⁻¹³ patient complaints,¹⁴ grievances,¹⁵ the level and seriousness of malpractice claims,^{16,17} or, perhaps most important, actual health and functional status outcomes¹⁸⁻²¹. Many researchers took both positions that patient perceptions have both inherent and instrumental value²¹⁻²⁴.

Quality of health care services in the hospital is the discrepancy between patient's perceptions of services offered by a particular hospital and their expectations about hospitals offering such services. The patient's expectations are derived from their perception of the ideal care standards of their previous experiences in the use of services²⁵. A health customer that is satisfied will be more likely to continue to use the services, spread positive views that will help healthcare providers get new patients without additional cost such as advertising²⁶.

Fundamentally, patients express their requirements in terms of what they need, want, prefer and demand with respect to the health care they receive²⁷. The requirements of patients could be considered as a demand for quality health care that health workers at primary level of health system attempt to meet²⁸⁻³⁰. Excellence in health care is what those in need of healthcare services at the primary health care level wish for, and it is also the main goal for those providing the care^{31,32}. Given that the combined product of health consumers' expectations and perceptions determined quality of health care services, this study therefore examined the expectations and perception the health consumers used in judging the quality of care services at the primary health care level.

Kwara State was created in 1967 as one of the 36 federating units of Nigeria. It is in the North central geopolitical zone of the country. It has an estimated population of 2,371,000, with annual growth rate of 3.4%. The Kwara State has 16 Local Government Areas grouped into 3 senatorial districts; the State capital is Ilorin³³. The public health service is organized into primary, secondary and tertiary health care levels. While the constitution is silent on the roles of the different levels of government in health services provision, the national health policy ascribes responsibilities for primary health care to local governments, secondary health care to states government and tertiary health care to the federal government³⁴. Primary health care (PHC), which forms the bedrock of the national health system, is the first level of care to the people³⁵.

METHODOLOGY

It was a community based cross sectional study using systematic sampling method to randomly selected 325 health consumers in selected primary health care facilities in Kwara State, Nigeria. Multistage sampling technique was adopted. First stage on the selection of primary health facility, all the Local Government Areas (LGAs) in the State were listed then all the primary health care facilities within the LGA in the State, from the listed PHC facilities in each LGA, two PHCs

facilities were randomly selected by balloting and total of thirty-two PHCs facilities were randomly selected by balloting for the study.

In second stage on the selection of health consumers, in each of the selected primary health care facility all the service delivery areas/units were listed. The list also includes supportive service area of the hospital and service delivery department / units include but not limited to hospital reception area, emergency services unit, general outpatient department, immunization unit, maternal and child health unit, laboratory services and health record unit, etc). In each of these service delivery areas/units, the first client i.e. health consumer to be registered and attended to by the health care providers was selected; thereafter every 2nd consecutive health consumers who had received any service or had at least one overnight stay and discharged in the hospital were randomly selected for interview in each service delivery point. However, health consumers that are in serious medical conditions that by virtue of illness will not be able to respond well to questions were excluded from study.

In the Measurements instruments used, the questions being asked comprises of open ended, dichotomous, rating and ranking questions. The tool was designed to elicit factual and definitive responses from the participants. The questionnaire was sectionalized into different parts of socio-demographic characteristics data, information on the respondents' perception and expectation of quality of health care services received. The research instrument was pre-tested among health consumers in other health institutions in Nigeria.

Research assistants made up of resident doctors, master students and medical students were recruited and trained for five days on data collection survey method. Proficiency of the interviewers was verified through role-play and pre testing. The researcher carried out visitation and checks in order to validate the collected data, each research assistant was recruited within the locality of the study site. Clearly define the outcome variables to be measured as products of the analysis, trained clerks entered the data from the interviews into the computer the same week the interviews are conducted and initially processed using excel then SPSS software version 20 packages were used for the analysis, *p* value, standard deviation and confident interval was calculated.

The qualitative section of the research was performed on one-to-one interviews with health consumers who had visited hospitals in the preceding six months especially women who delivered at the health facility. Three health consumers were interviewed in each PHC facility. Topics discussed include socio-cultural perception and expectation of quality health care services, feeling and if there any difference between

the ways clients are treated. After each session, the tape recordings was transcribed and reconciled with the notes taken. The full recording was then edited to remove responses / session that was poorly transcribed. Finally, similar responses were group-coded, analyzed and presented.

After the ethical clearance from University of Ilorin ethical review committee, a letter of introduction was obtained from the department of epidemiology and community health, university of Ilorin. This was presented to the relevant authorities of the respective health Institutions that was selected requesting for their consent and participation in the study. During the surveys, consent was obtained from the respondents. Informed sheets and consent forms were translated into relevant languages. Verbal consent was obtained prior to the survey, and all verbal consent was witnessed by a field worker. All survey data was kept confidential. At each facility verbal consent was obtained to avoid suspicious of having to sign documents that may be used to identify them or their outlet/facility. Participants were allowed to drop out at any time during the survey and do not have to respond to questions they do not wish to answer. Ethical approval received from institution and protocol Identification code is UERC/CLS/007.

RESULTS

This study response rate was 100%, there were more females 300 (92.3%) than male respondents, the predominant religious groups are Christianity and Islam accounting for 75 (23.1%) and 250(76.9%) respectively. Married respondents are 175(53.8%) while 150(46.2%) were single. On educational status post- secondary education 175(53.8%) while others had secondary education 100(30.8%). Only 125 (38.5%) of the respondents had monthly income of ≤ \$28.00 (Table I)

As seen in (Table II), explained positive perception 88 (27.1) and patient expectation 178(54.7) of total variance for medical doctor satisfaction scores. The clients had high positive expectation 223(68.5%) of the health facility premises but not much in difference on positive perception (61.6%) and negative expectation (61.0%) of diagnostic services. Other domains disparity in positivity or negative expectations or perceptions in services quality the clients received from nurses, during admission to the hospital, the meals giving to them with services received in the wards in Table II. The reliability of the existence of quality gap in term of overall perception (5.80) and expectations (6.57) of services is showed in Table III.

An in-depth interview analysis of the tool showed that most of the patients were positive, such as "the service was good and doctors and nurses responded

to questions quickly and completely". Though, prolong waiting time, unpleasant smelling of the reception area and young age of some nurses was an issues. In Table IV, the chi square test on gender showed no disparity in overall expectation-perception, though females had higher expectation than the male counterparts. Correlation test revealed significant changes in socio-demographic variables such as age,

TABLE I: SOCIODEMOGRAPHIC CHARACTERISTIC OF THE HEALTH CONSUMERS AT MPHCH

Variable/ Factors	MPHC (N=325)	Frequency (%)
Age groups	≤ 30	111 (34.2) **
	31- 40	100(30.8)
	41- 50	56 (17.2)
	51- 60	44 (13.5)
	≥ 61	14 (4.3)
Gender	Male	25 (7.7) **
	Female	300 (92.3)
Religion	Islam	250 (76.9) **
	Christianity	75 (23.1)
	Traditional	0(0)
Education Status	No Formal Edu.	00 (00.0) **
	Primary	50 (15.4)
	Secondary	100 (30.8)
	Post-Sec.	175 (53.8)
Marital Status	Single	150 (46.2) **
	Married	175 (53.8)
	Separated	0 (0)
Occupation	Civil Servant	75 (23.1) **
	Self Employed	50(15.4)
	Unemployed	80(24.6)
	Farmers	70 (21.5)
	Artisans	50(15.4)
	Students	0(0.0)
	Other	0(0.0)
Monthly Income (\$)	≤28.00	125(38.5)*
	28.00 – 139.00	125 (38.5)
	139.00 – 270.00	50 (15.4)
	≥270.00	25 (7.6)
Overall	Total	325(100)

* $p < 0.05$; ** $p < 0.01$; + $p = 0.047$; # $p = 0.05$
 *variables significant for levels of care are highlighted
 Mean age of 35.95±10.5

TABLE II: DOMAINS OF QUALITY OF HEALTH CARE SERVICES HEALTH CONSUMERS EXPECTATION - PERCEPTION OF SERVICES RECEIVED

S/N	Domains variables	Expectation Positive Frequency (%)	Expectation Negative Frequency (%)	Perception Positive Frequency (%)	Perception Negative Frequency (%)
1	Hospital Premises and Employees	223(68.5)	102(31.5)	71(21.7)	254(78.3)
2	Quality of Services from Medical Doctor	178(54.7)	147(45.3)	88(27.1)	237(72.9)
3	Quality of Diagnostic Services	125(38.4)	200(61.6)	198(61.0)	127(39.0)
4	Quality of Services from Nurses	205(63.6)	120(36.4)	4(14.0)	321(86.0)
5	Quality of Admission Services	316(97.3)	9(2.7)	37 (11.1)	288(88.9)
6	Quality of Meals & Laundry (ML) Services	106 (32.6)	219(67.4)	215(66.4)	110(33.6)
7	House Keeping Services Clinics, and Wards	177(54.3)	148 (45.7)	24(7.2)	301(92.8)

TABLE III: RELIABILITY AND PAIRED T TEST

Variables	Mean score	SD	t-test	Cronbach's alpha
Expectation	6.57	0.234	1.08**	0.947
Perception	5.80	0.135		0.858

Note * means significant at 0.05 level

TABLE IV: RELATIONSHIP BETWEEN SOCIO-DEMOGRAPHIC CHARACTERISTICS AND PATIENTS EXPECTATION- PERCEPTION ON QUALITY OF HEALTH CARE SERVICES

Socio-demographic Variables		PHCF (%) N=325 Expectation Freq.(%) Perception Freq. (%)	
Age Group (years)	≤ 30	100 (36.4)	25 (50.0)
	31- 40	100 (36.4)	0 (0.0)
	41- 50	75 (36.4)	25 (50.0)
	51- 60	0 (0.0)	0 (0.0)
	≥ 61	0 (0.0)	0 (0.0)
	χ²/p-v	27.330/ <0.001	
Gender	Male	25 (9.1)	0 (0)
	Female	250(90.9)	50 (100.0)
	χ²/p-v	4.924/0.026	
Religion	Islam	200(72.7)	50 (100.0)
	Christian	75 (27.3)	0 (0)
	Other	0 (0)	0 (0)
	χ²/p-v	17.727/<0.001	
Education status	None	0 (0)	50 (100.0)
	Primary	50 (18.2)	0 (0)
	Secondary	50 (18.2)	0 (0)
	Tertiary	175(63.6)	0 (0)
	χ²/p-v	32.50/<0.001	
Marital Status	Single	100(36.4)	50 (100.0)
	Married	17 (63.6)	0 (0)
	Separated		
	χ²/p-v	68.939/<0.001	
Occupation	Civil.Servt	175(63.6)	0 (0)
	Self Emp	0 (0)	0 (0)
	Unemploy	75 (27.3)	0 (0)
	Farmer	25 (9.1)	50 (100.0)
	Artisan	0 (0)	0 (0)
	Students	0 (0)	0 (0)
	Other	0 (0)	0 (0)
	χ²/p-v	1.974/<0.001	

Monthly IncomeX000	≤10	50 (22.2)	0 (0)
	11 – 50	100(44.4)	25(100.0)
	501 – 100	50 (22.2)	0 (0)
	≥101	25 (11.1)	0 (0)
	$\chi^2/p-v$	27.778/<0.001	
In-Outpatient	Inpatient	125(45.5)	0 (0)
	Outpatient	150(54.5)	50(100.0)
	$\chi^2/p-v$	36.932/<0.001	

occupation, educational status and services quality variable examined. With increase in level of education there is decrease in perception scores of the clients ($p_v = <0.001$).

DISCUSSION

Hospitals establishments like many other organizations are playing an increasingly important role that affects the health, wellbeing and economy of less-developed countries. One of the strong points of this study is the high percentage of responses, which reduces the non-response bias. This response rate was slightly higher than 99.9% rate of similar research in Latin American,^{36,37} and higher than the 63% in a study conducted among six hospital wards at the Academic medical center, Amsterdam.

With mean age of 35.95±10.5, it implied that respondent's experiences are still recent enough for accurate recall and that majority of the respondents are within the reproductive age group (women of child bearing age). The pattern of age distribution of health consumers is in keeping with the population pyramid of most developing countries with a preponderance of the mid-aged compared with few elderly individuals WHO technical report series,³⁸ there are more females than males this could be due to the fact that health care received at the primary health care in Nigeria was supported by midwives' services scheme programme of federal government as the cost of delivery was relatively free. Though, majority of the respondents are married which is keeping with national demographic health survey³⁹. However, significant difference observed in expectation and perception gaps in patients age, occupation and income level, the results was different from study conducted among patients with refractive visual disorders⁴⁰.

The higher proportion of Muslim respondents in this survey is not surprising because large percentages of people in study area are Muslims. The religious affiliation of the respondents is important since some beliefs about sickness and getting well were premised on the socio-cultural and religious practices of the individuals. Though, differ from other study,⁴⁰ on

traditional and religious practices affecting the health of other women in Nigeria that many Muslims women are kept in purdah(Vail), during their reproductive years with social restrictions and low utilization of primary health care services in the hospital, utilization of primary health care services by many Muslim women may be supported with the fact that all the respondents interviewed were in one form of occupation or the other (Table I). More than half of respondents had formal education which is higher than the national figure of 43%. Educational level is an important index of health status, wellbeing and provided greater access to health information⁴¹.

Though high expectation-perception gap was found in this study, however, the use of only perceptions to determine quality of healthcare services was found in others studies^{42,43} conducted among nurses and dentists respectively, where it was reported that addition of expectations is unnecessary especially in the hospitals setting where expectations from healthcare providers usually range from high to very high due to the importance of all different dimension of the services provided in the hospital. In another study⁴⁴ on a gap analysis of professional on quality of services, supported that use of only perceptions for determination of quality of services as it reveled recent events, problems or challenges in the establishment on services provided to the consumers which will enhance immediate short-term solution than to start on long-term planning. However, this study was corroborated by study on customers' perception and expectation of primary health care services quality in health centers of Bandar Abbas⁴⁵ found that periodic (e.g. quarterly, biannual or annually) collection of data on consumers' expectation can provide an insight for the long-term planning and improvement process for the hospital.

The results of comparing individuals' expectation-perception with the socio-demographic characteristics shown significant effects (Table IV). For example, classification of educational level had significant effect on expectation-perception of clients at the significant level of $p < 0.05$ which may be due to the fact that health consumers with lower educational status had lower expectations of quality of healthcare services than the health consumers with tertiary education level. This study also found that women's expectation of quality of healthcare services was higher than that of the men expectation of quality of healthcare services. The women's higher expectations compared with the men had been reported in a study conducted on quality of screening service for chronic kidney-disease research evidence from Taichung City⁴⁶.

Apart from socio-demographic characteristics effects on patients' expectation- perception of quality of

healthcare, other study have shown that patients 'expectations of quality of healthcare exceed perceptions of the healthcare as the patients felt that waiting time of more than an hour to receive the service was excessive and that the healthcare providers did not respond fast enough when problems arose in the hospitals⁴⁷. It was also found that expectation-perception gap enables healthcare providers to analyze the important of health services from the consumer perspective and to prioritized the areas or sections in the hospital that need immediate improvement for clients. Expectation-perception gap analysis of different domains in the hospital ensured proper allocation of resources starting with areas or units deemed most important for a short term than for long term.

The results of this research showed increase expectations of health consumer's in hospital premises and employees. Expectation from medical doctors and nurses working in the hospital at the primary health care level, quality of diagnostic services received from time of admission to wards and in the clinics. Overall respondent's expectation on quality of health care services received was more than average while perception of the services was less than one third. When compared with other studies conducted among patients in district hospitals and primary health centers in India⁴⁸ the scores were less than the findings in this study.

There were extremely gaps in housekeeping services clinics and wards, nurses, and environmental premises. The reliability, assurance, and empathy were affected by the gaps. This study suggested that health workers should increase discussions and communication with their patients for satisfaction of services received as reported in other study⁴⁹.

CONCLUSIONS

This study found that expectation- perception is a major determinant of quality of health care services at primary health care levels; there is disparity in expectation-perception gap among males and females respondents. However, significant relationship occurred in patients' age, occupation, educational status and overall expectation -perception of quality care received. Behaviour variables and the methodology of this stud can be easily adopted by the research team in the hospital to determine patient behaviours with service quality expectation.

REFERENCES

1. Leebov W. The quality quest: a briefing for health-care professionals. *J HealthCare Quality*. 1991; 15(4): 42-47.
2. Sun BC, Adams J, Orav EJ, Rucker DW, Brennan TA, Burstin HR. Determination of patients satisfaction and willingness to return with Emergency care. *Ann Emerg Med*. 2000; 35(5):426-34.
3. Hoy E. Measuring patient experiences of care. *Bull Am Coll Surg*. 2008; 93(5):13-6.
4. Farooqi JH. Patient expectation of general practitioner care. *Midd East J Fam Med*. 2005; 3 (3): 6-9.
5. Coulter A. After Bristol: putting patients at the centre. *BMJ Quality & Safety*. 2002;11(2):186-88.
6. Gerteis M, Edman-Levitan S, Daley J, Delbanco TL, editors. *Through the Patient's Eyes: Understanding and Promoting Patient-Centered Care (First Edition)*. Jossey-Bass, San Francisco. 1993.
7. Lauver DR, Ward SE, Heidrich SM, Keller ML, Bowers BJ, Brennan PF, et al. Patient centered interventions. *Res Nurs Health*. 2002; 25(4): 246-55.
8. Dwamena F, Holmes-Rovner M, Gaulden CM, Jorgenson S, Sadigh G, Sikorskii A, et al. Interventions for providers to promote a patient-centred approach in clinical consultations. *Cochrane Database Syst Rev*. 2012; 12:CD003267. doi:10.1002/14651858.CD003267.pub2.
9. McLaughlin CP, Kaluzny AD. Building client centered systems of care: choosing a process direction for the next century. *Health Care Manage Rev*. 2000; 25(1):73-82.
10. Mead N, Bower P. Patient centredness: A conceptual framework and review of the empirical literature. *Soc Sci Med*. 2000; 51(7):1087-1110.
11. Bartlett JA. Addressing the challenges of adherence. *J Acquir Immune Defic Syndr*. 2002; 29 suppl 1:S2-S10.
12. Brown R. Behavioral issues in asthma management. *Pediatr Pulmonol Suppl*. 2001; 21:26-30
13. Golin CE, DiMatteo MR, Gelberg L. The role of patient participation in the doctor visit. Implications for adherence to diabetes care. *Diabetes Care*. 1996; 19(10):1153-64
14. Taylor DM, Wolfe R, Cameron PA. Complaints from emergency department patients largely result from treatment and communication problems. *Emerg Med (Fremantale)*. 2002; 14(1):43-49.
15. Gruman JC. An expanded view of health: implications for how healthcare works. *Behav Med*. 1994; 20(3):119-22.
16. Hickson GB, Clayton EW, Githens PB, Sloan FA. Factors that prompted families to file medical malpractice claims following perinatal injuries. *JAMA* 1992; 267(10): 1359-63.
17. Hickson GB, Federspiel CF, Pichert JW, Miller CS, Gauld-Jaeger J, Bost P. Patient complaints and malpractice risk. *JAMA*. 2002; 287(22):

- 2951-57.
18. Cleary PD, McNeil BJ. Patient satisfaction as an indicator of quality care. *Inquiry*. 1988; 25(1): 25-36
 19. Covinsky KE, Rosenthal GE, Chren M, Justice AC, Fortinsky RH, Palmer R, et al. 1998. The relation between health status changes and patient satisfaction in older hospitalized medical patients. *J Gen Intern Med*. 1998; 13:223-29
 20. Fremont AM, Cleary PD, Hargraves JL, Rowe RM, Jacobson NB, Ayanian JZ. Patient-centered processes of care and long-term outcomes of myocardial infarction. *J Gen Intern Med*. 2001; 16 (12):800-8.
 21. Jung H, Wensing M, Olesen F, Grol R. Comparison of patients' and general practitioners' evaluations of general practice care. *Qual Health Care*. 2002; 11(4):315-19
 22. Stewart M, Meredith L, Brown JB, Galajda J. The influence of older patient physician communication on health and health-related outcomes. *Clin Geriatr Med*. 2000; 16(1): 25-36.
 23. Maly RC, Bourque LB, Engelhardt RF. A randomized controlled trial of facilitating information giving to patients with chronic medical conditions: effects on outcomes of care. *J Fam Pract*. 1999; 48(5): 356-63.
 24. Marshall GN, Hays RD, Mazel R. Health status and satisfaction with health care: results from the medical outcomes study. *J Consult Clin Psychol*. 1996; 64(2):380-90.
 25. Kucukarslan SN, Nadkarni A. Evaluating medication-related services in a hospital setting using the disconfirmation of expectations model of satisfaction. *Res Soc Adm Pharm*. 2008; 4(1): 12-22. doi: 10.1016/j.sapharm.2007.01.001.
 26. Zeithaml VA, Bitner MJ, Gremler DD, eds. *Services Marketing (4th International Edition)*. In: McGraw-Hill, Inc, New York, NY, 2005.
 27. Reck DL. Can and should nurses be aware of patients' expectations for their nursing care? *Nurs Adm Q*. 2013; 37(2):109-15. doi: 10.1097/NAQ.0b013e3182869e03.
 28. Courtenay M, Carey N, Stenner K, Lawton S, Peters J. Patients' views of nurse prescribing: effects on care, concordance and medicine taking. *Br J Dermatol*. 2011; 164(2):396-401. doi: 10.1111/j.1365-2133.2010.10119.x.
 29. Mohamed HA, Sayed HY. Patients' perceptions as indicators of quality of nursing service provided At Al Noor Specialist Hospital at Makkah Al Moukarramah, KSA. *J Am Sci*. 2013; 9(5):71-8.
 30. Shaller D. *Patient-Centered Care: What does it take?* New York: Commonwealth Fund; 2007.
 31. Abrahamsen Grondahl V. Patients' perceptions of actual care conditions and patient satisfaction with care quality in hospital (Phd dissertation). Karlstad: Karlstads Universitet; 2012.
 32. Gerense H, Solomon K, Birhane M, Medhin BG, Mariam TH, Guesh Kibra, et al. Quality of nursing care among in-patient of medical-surgical ward in Axum St. Marry Hospital, Tigray, Ethiopia 2015. *Enz Eng*. 2015; 4:132. doi:10.4172/2329-6674.1000132.
 33. National Planning Commission. *Children's and women's rights in Nigeria: A wake-up call. Situation Assessment and Analysis*. 2001. ISBN No. 92-9186-021.
 34. Saka MJ, Saka AO, Isiaka SB, Agbana BE, Bako IA. Health related policy reform in Nigeria; - Empirical analysis from 2001 to 2010; The Past, Trend and Future Directions for Improved Sustainable Health Financing and Development. *Int J Acad Res Bus Soc Sci*. 2012; 2(3); 27-37.
 35. Saka MJ, Saka AO, Oloriegbe IY. Overview of legal Implication and Health Consequences of Human Trafficking in Nigeria. *Ilorin J Education*. 2017; 36:1-8.
 36. Ndhlovu L. *Quality of Care in Family Planning Service Delivery in Kenya: Clients' and Providers' Perspectives*. Nairobi, Kenya: Population Council, Africa Operation Research and Technical Assistance Project. 1995; 1-84.
 37. Bankauskaite V, Saarelma O. Why are people dissatisfied with medical care services in Lithuania? A qualitative study using responses to open-ended questions. *Int J Qual Health Care*. 2003; 15(1):23-29.
 38. World Health Organization. *Report of a WHO Expert Committee on Health of the Elderly. Demography and Public Health Aspects of Population Ageing*. WHO Tech Rep Ser. 1989; 779:14-17.
 39. Lin DJ, Sheu IC, Pai JY, Bair A, Hung CY, Yeh YH, et al. Measuring patient's expectation and the perception of quality in LASIK services. *Health Qual Life Outcomes*. 2013; 7: 63-67. doi: 10.1186/1477-7525-7-63.
 40. Johnson BC. Traditional practices affecting the health of women in Nigeria. In: Baasher T, Bannerman RH, Rushwan H, Sharaf I, editors. *Traditional Practices Affecting the Health of Women and Children*. Alexandria, Egypt: World Health Organization. 1982: 23-25.
 41. Saka MJ, Abdul FI, Saka AO, Odusolu PO, Okesina BS, Bako IA, et al. Assessment of Infrastructure and Community Supply in Nigeria Private Health Facilities: Implication on the implementation for Maternal and Child health Care Policies. *Trop J Health Sci*. 2012; 19(1):31-34.

42. John J. Patient Satisfaction: The Impact of Past Experience. *J Health Care Mark.* 1992; 12:(3):56-64.
43. Paul D. What is the "Best" approach for measuring service quality of Periodontists? *Clinical Research Regulatory Affairs.* 2003; 20(4): 457-468.
44. Brown SW, Swartz TA. A Gap Analysis of Professional Service Quality. *Journal of Marketing.* 1989; 53(2):92-98.
45. Agha Molaei T, Zare SH, Poudat A, Kebriaei A. Customers Perception and Expectation of Primary Health Care Services quality in Health Centers of Bandar Abbas. *Med J Hormozgan Uni.* 2007; 11 (3):173-79.
46. Lin DJ, Li YH, Pai JY, Sheu IC, Glen R, Chou MJ, et al. Chronic Kidney-Disease screening service quality: Questionnaire survey research evidence from Taichung City. *BMC Health Serv Res.* 2009; 9: 239. doi: 10.1186/1472-6963-9-239.
47. Suki N M, Lian JC, Suki NM. Do patients 'perceptions exceed their expectations in private healthcare settings? *Int J Health Care Qual Assur.* 2011; 24(1):42-56.
48. Lin HC, Xirasagar S, Laditka JN. Patient perceptions of service quality in group versus solo practice clinics. *Int J Qual Health Care.* 2004; 16 (6):437-45.
49. Chen JY, Tao ML, Tisnado D, Malin J, Ko C, Timmer M, et al. Impact of Physician-Patient Discussions on Patient Satisfaction. *Medical Care.* 2008; 46(11):1157-62. doi: 10.1097/MLR.0b013e31817924bc.



AUTHOR AFFILIATION:

Dr. Saka Mohammed Jimoh (*Corresponding Author*)

Epidemiology and Community Health Department
Faculty of Clinical Sciences, College of Health Sciences
University of Ilorin, Ilorin Nigeria.
Email: sakamj1@yahoo.com, sakamj@unilorin.edu.ng

Dr. Akande Tanimola Mankanjuola

Epidemiology and Community Health Department
Faculty of Clinical Sciences, College of Health Sciences
University of Ilorin, Ilorin Nigeria.

Dr. Saka Aishat Oluwatoyin

Department of Pediatric and Child Health
Faculty of Clinical Sciences, College of Health Sciences
University of Ilorin, Ilorin Nigeria.

Dr. Bello Fatai A

Department of Community Health
College of Health Sciences
University of Ilorin, Ilorin Nigeria.

Isiaka Sulu Babaita

Department of Business Administration
Faculty of Management Sciences
University of Ilorin, Ilorin Nigeria.