

Evaluation of Pattern and Impact of Electrolytes Abnormalities in Critically Ill Covid-19 Patients

Anita Haroon, Syed Ali Abbas, Amanullah Khan, Momina Ali, Rija Qazi, Ajeet Kumar

ABSTRACT

OBJECTIVE: To evaluate the pattern of serum electrolytes abnormalities and their impact on ICU admitted Covid-19 patient outcomes.

METHODOLOGY: This retrospective study was carried out at OMI hospital and Dr. Ziauddin Hospital, Karachi, Pakistan, between August to December 2020. Total 102 PCR positive, ICU admitted with severe Covid-19 patients as per WHO criteria were included. The patient's demographic characteristics, clinical features including co-morbidities, electrolytes reports at the time of admission, length of ICU and/or hospital stay, and outcome (expired/survived) were evaluated.

RESULTS: Biochemical testing found abnormal electrolyte levels in 90.2% ICU admitted Covid-19 patients. Electrolytes abnormalities including hyponatremia 45.1%, hypermagnesemia 40.2%, hypocalcemia 31.4%, hyperchloremia 23.5% and hyperphosphatemia in 20.6% patients. Out of the total, 28.4% of patients needed invasive respiratory support, and 37.3% could not survive. A higher incidence of mortality (39.1% vs. 20%) was seen in patients with electrolytes abnormalities compared to those presented with normal values.

CONCLUSION: Electrolyte abnormalities were found in 90% of the ICU Admitted Covid-19 patients. The most common abnormalities found among the patients were hyponatremia, hypermagnesemia, and Hypocalcemia. The findings revealed that several electrolyte imbalances harm patients' in-hospital outcomes. Electrolyte assessment of Covid-19 patients at the time of admission would be helpful in risk stratification for adverse outcomes.

KEYWORDS: Covid-19, critically ill, ICU stay, serum electrolytes, abnormalities, respiratory support, death.

This article may be cited as: Haroon A, Abbas SA, Khan A, Ali M, Qazi R, Kumar A. Evaluation of Pattern and Impact of Electrolytes Abnormalities in Critically Ill Covid-19 Patients. J Liaquat Uni Med Health Sci. 2022;21(01):16-22. doi: 10.22442/jlumhs.2022.00905. Epub 2022 January 27.

INTRODUCTION

Within a few months of the World Health Organization (WHO) announcing a global pandemic, several million cases of coronavirus disease (Covid-19) have been reported globally since December 2019¹. Different updates on the clinical characteristics of infected patients have been unveiled since Covid-19, an infectious disease caused by severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2), emerged in Wuhan, China². With multisystem involvement, Covid-19 is a potentially fatal disease. However, the human respiratory system is its primary target, which triggers severe pneumonia and develops acute respiratory distress syndrome (ARDS) in the respiratory system. In addition, the condition can affect multiple organs system of the human body, including the neurological, renal, gastrointestinal and coagulation systems³. An increase in morbidity and mortality among critically ill patients is associated with electrolyte imbalances (sodium, calcium, potassium, chloride, phosphate, and magnesium)⁴. Electrolyte imbalances in patients were stated in several recently conducted studies. Electrolyte imbalances have contributed to cardiovascular and renal diseases, increased hospitalization time, and

mortality caused by SARS-CoV-2 infection^{5,6}. Some data has been presented in early Covid-19 studies that electrolyte abnormalities, sodium, calcium, potassium, and chloride, can also be found upon patient presentation⁶⁻⁸.

By binding Angiotensin-converting enzyme 2 (ACE2) to the cell membrane, human cells invade by SARS-CoV-2. In several human tissues where ACE2 is widely distributed, e.g., lungs, heart, liver, and kidney, organ dysfunction is seen. For the renin-angiotensin system (RAS) main axis, ACE2 is the primary counter-regulatory pathway, which plays a vital role in regulating blood pressure and electrolytes. This virus binds to ACE2 and increases ACE2 degradation, thus reducing the RAS counteract of ACE2. The result is the rise of serum sodium level and water re-absorption, subsequently high blood pressure and potassium (K⁺) excretion. In addition, Covid-19 patients may also have GI symptoms, such as vomiting and diarrhea. The effects of Covid-19 on RAS and the gastrointestinal system are likely to lead to disturbances of electrolyte homeostasis and pH^{2,9}. Previous studies also indicated such imbalances of serum electrolyte and glucose levels in Covid-19 individuals^{6,10,11}.

There is an association between hyponatremia and atypical pneumonia, which leads to ICU admission, longer duration of stay, higher hospital costs, and high incidence of mortality². Duan J et al.¹² reported the transition of the Covid-19 to severe stage. These markers, sodium, potassium, and chloride levels, had high predictive capacity. Low baseline sodium, chloride, and calcium levels were found by Tezcan ME 2020³ to be correlated with higher mortality rates, increased Intensive care unit stay and mechanical ventilator requirements, and prolonged hospital stay. The predominant electrolyte abnormality associated with adverse findings is hyponatremia². Low sodium, potassium, and calcium serum concentrations are directly related to the severity of the disease. Though it is not yet clear if there is a risk of Covid-19-related hyponatremia, Hypocalcemia, or Hypocalcemia, more clinical knowledge will be required to interpret these abnormalities. These conditions, in particular hypokalemia, may have considerable consequences for the treatment of patients and may theoretically lead to the unraveling of the fundamental pathogenic pathways of Covid-19. Acute respiratory distress syndrome (ARDS) and acute cardiac injury, which are frequent complications of covid-19, intensify hypokalemia, especially in patients with underlying lung or heart disease^{6,13,14}.

Poor prognosis and severe disease have been related to a low potassium level at enrollment, but little is understood about the dynamics over time. In previous research, chloride homeostasis has been briefly addressed without obtaining any prognostic benefit at admission. The prevalence of electrolyte abnormalities is significantly higher among severely ill Covid-19 patients. Serum sodium (Na⁺) level in 57-77.1%, potassium (K⁺) level in 50%, calcium (Ca²⁺) level in 28.6%, magnesium(Mg) level in 15.7% patients were found to be low and high magnesium level was found in 7.14% Covid-19 patients^{2,6,15,16}.

Our research aims to describe the dynamics of serum electrolytes at presentation and the outcomes of patients with Covid-19 who are critically ill. This study aims to help ICU health providers optimize Covid-19 patients' electrolyte management. At present, there is limited evidence about severe Covid-19 patients. Therefore, knowledge of these corona virus-induced metabolic abnormalities will play a role in disease prognosis, initial therapeutic measures, and disease surveillance. Knowledge and understanding of the possible pathophysiological pathways of Covid-19 could provide an opportunity to develop a new therapeutic strategy. Therefore, we tried to strengthen our present understanding of electrolyte imbalance in ICU admitted Covid-19 patients. The current study with structured measurements of serum levels with clinical evaluation of electrolytes input and output will add significant value to the existing literature. It will also provide crucial information for decision-making in

the current public health crisis.

METHODOLOGY

After taking ethical approval, this retrospective study was conducted at OMI hospital and Dr. Ziauddin Hospital, Karachi. The sample size was calculated using the online sample size calculator Open Epi. Taking statistics of hypomagnesemia 7.14% in Covid-19 patients admitted in ICU², at the margin of error 5% and confidence interval 95%. Data of 102 Covid-19 patients were obtained using the "Non-probability consecutive sampling technique" from hospitals' medical records. Only those Covid-19 adult (>18 years age) patients irrespective of gender who were treated at the intensive care unit (ICU) from August to December 2020 were included.

Covid-19 or SARS-CoV-2 was defined as a positive result of real-time reverse transcriptase-polymerase chain reaction (PCR) assay of nasal and pharyngeal swabs and following WHO¹⁷ criteria for only severe (signs of pneumonia with respiratory rate \geq 30 breaths/min or SpO₂ < 90% on room air) and critical (development of Acute Respiratory Distress Syndrome, septic shock or multi-organ dysfunction) Severe Covid-19 patients admitted in ICU were enrolled in this study. Patients less than 18 years of age, who had other causes of electrolyte imbalances, chronic organ failure, immunodeficiency, and terminal cancer, were excluded.

Electrolytes levels were obtained from the laboratory database. The patient's electrolyte values, i.e., serum sodium, potassium, chloride, magnesium, calcium, bicarbonate, phosphorus, and albumin at the time of admission, were extracted from their medical records. According to the standard laboratory reference range, the values of these markers are further categorized as hypo and hyper.

Normal reference values of different electrolytes are as followed: 135-145 mmol/L – Serum sodium level, 3.5-5 mmol/L- Potassium level, 95-105 mmol/L- Chloride level, 1.7-2.2 mg/dL- Magnesium level, 8.5-10.2 mg/dL – calcium level, 2.5 – 4.5 mg/L-phosphorus level. Demographic and clinical characteristics, including age, gender, co-morbid status, were also obtained from patients' medical records. Patient outcomes (Noninvasive/invasive reparatory support, duration of hospitalization, duration of ICU stay, and in-hospital mortality) and the impact of the electrolyte abnormalities on outcomes were also evaluated in this study.

Data was entered and analyzed by using SPSS version 24. Descriptive statistics were calculated for study variables. Mean and standard deviation were computed for quantitative variables, and frequency and percentages were calculated for qualitative variables. Associations between outcomes and confounders were observed using the Chi-square/ Fisher exact test. Comparisons of quantitative

variables were made using an independent t-test. At P-value ≤ 0.05 as significant and power of the test kept 80%.

RESULTS

This study included 102 critically ill Covid-19 patients admitted in ICUs at two different private hospitals in Karachi, Pakistan. In this cohort mean age of the patients was 63.2+/-13.2 years. 57.8% were over 60 years of age, 37.3% were between 41-60 years and remained below 40 and above 18. Male patients were 56.9%. Hypertension, diabetes mellitus, and IHD were most frequently found co-morbidities (Table I). Electrolyte levels at admission were assessed and found abnormal in 90.2% Covid-19 patients. Where only 1 abnormal level was found in 22(21.6%), 2 in 30 (29.4%), 3 in 20(19.6%), 4 in 16(15.7%), 5 in 3(2.9%) and 6 abnormal electrolytes found in 1(1%) patients. Approximately 70% of patients had more than one electrolyte imbalance. 64.7% patients had hyponatremia, 40.2% hypermagnesemia, 36.3%, hypocalcemia, 31.4%, 23.5% hyperchloremia, and 20.6% were found to have hyperphosphatemia. The most common abnormalities found in our study were hyponatremia, hypermagnesemia, and Hypocalcemia, while less frequent abnormalities were hypokalemia, hyperkalemia, hypochloremia, hypomagnesemia, and

hypophosphatemia. (Figure I) 71.6% of patients required noninvasive respiratory support, and the remaining 28.4% needed invasive mechanical ventilators. The mean ICU stay of Covid-19 patients was 7.4+/-5.8 days, while overall hospital stay was 10.1+/-6.4 days. Furthermore, 37.3% of patients died. (Table I)


A comparison of overall electrolyte abnormalities with demographic, clinical profile, and outcome of the patients was made. Most of the factors had no statistical difference (P-value > 0.05), while gender and respiratory outcome support had a significant association with electrolyte abnormality (P-value<0.05). (Table I)

A comparison of electrolyte abnormalities with the outcome of the patients was made. Although no statistical difference was found (P-value>0.05), the frequencies showed a high prevalence of electrolyte abnormalities in patients who died. Among patients who died, 31.6% had Hypocalcemia, 44.7% had hyponatremia. Respiratory support showed a significant association with patient outcomes. Out of all patients who needed invasive mechanical ventilation, only 20.3% survived. On the other hand, 79.7% of those on noninvasive ventilation could survive. ICU and hospital stay was relatively higher in patients who had electrolyte abnormalities and who

TABLE I: DISTRIBUTION AND COMPARISON OF DEMOGRAPHIC, DISEASE-RELATED PROFILE AND OUTCOMES OF THE COVID-19 PATIENTS

Associated factors	Electrolytes Abnormalities		Total	P-values	
	Yes	No			
Age groups	>40 years	5(5.4%)	0(0%)	5(4.9%)	0.260**
	41-60 years	32(34.8%)	6(60%)	38(37.3%)	
	>60 years	55(59.8%)	4(40%)	59(57.8%)	
	Mean+/-SD			63.2+/-13.2	
Gender	Male	49(53.3%)	9(90%)	58(56.9%)	0.026*
	Female	43(46.7%)	1(10%)	44(43.1%)	
Co-morbidities	Diabetes mellitus	54(58.7%)	3(30%)	57(55.9%)	0.083**
	Hypertension	54(58.7%)	4(40%)	58(56.9%)	0.257**
	Asthma	4(4.3%)	1(10%)	5(4.9%)	0.432**
	COPD	4(4.3%)	0(0%)	4(3.9%)	0.501**
	IHD	25(27.2%)	3(30%)	28(27.5%)	0.849**
Respiratory support	Invasive	29(31.5%)	0(0%)	29(28.4%)	0.029*
	Non-invasive	63(68.5%)	10(100%)	73(71.6%)	
ICU stay		7.67+/-5.94	5.10+/-4.46	7.4+/-5.8	0.188**
Hospital stay		10.39+/-5.6	7.90+/-4.84	10.1+/-6.4	0.248**
Outcome	Expired	36(39.1%)	2(20%)	38(37.3%)	0.235**
	Survive	56(60.9%)	8(80%)	64(62.7%)	
Total		92(90.2%)	10(9.8%)	102(100%)	

FIGURE I: PATTERN OF ELECTROLYTES ABNORMALITIES



died compared to the patients who survived. While there was no difference observed in-hospital stay among patients who died or survived.

A comparison of electrolyte abnormalities with respiratory support received by the patients was made. Similar to the previous finding, no statistical difference was found here as well (P-value > 0.05), but the frequencies showed there is a relatively high prevalence of electrolyte abnormalities in patients who were on invasive support. Among those on invasive support, 58.6% had hyponatremia, 48.3% had hypermagnesium, and 37.9 had Hypocalcemia. ICU and hospital stay was higher in patients who were on invasive respiratory support than patients who were on noninvasive support. (Table II)

This study also explored the association between

TABLE II: COMPARISON OF ELECTROLYTE ABNORMALITIES AND OUTCOMES IN COVID-19 PATIENTS

Electrolyte abnormalities	Respiratory distress		Sig.	Outcome		Sig.
	Noninvasive	Invasive		Expired	Survived	
Hyponatremia	29(39.7 %)	17(58.6%)	0.084	17(44.7%)	29(45.3%)	0.955
Hypokalemia	13(17.8%)	5(17.2%)	0.946	5(13.2%)	13(20.3%)	0.359
Hyperkalemia	9(12.3%)	4(13.8%)	0.537	6(15.8%)	7(10.9%)	0.545
Hypochloremia	6(8.2%)	2(6.9%)	0.823	5(13.2%)	3(4.7%)	0.145
Hyperchloremia	18(24.7%)	6(20.7%)	0.670	11(28.9%)	13(20.3%)	0.320
Hypomagnesaemia	8(11%)	5(17.2%)	0.290	3(7.9%)	10(15.6%)	0.207
Hypermagnesaemia	27(37%)	14(48.3%)	0.294	18(47.4%)	23(35.9%)	0.255
Hypocalcaemia	21(28.8%)	11(37.9%)	0.368	12(31.6%)	20(31.2%)	0.972
Hyperphosphatemia	18(24.7%)	3(10.3%)	0.107	11(28.9%)	10(15.6%)	0.108
Hypophosphatemia	7(9.6%)	3(10.3%)	0.582	4(10.5%)	6(9.4%)	0.522
Outcomes						
ICU Stay	6.7+/-5.7	9.1+/-6.0	0.576	8.4+/-6.0	6.8+/-5.7	0.185
Hospital Stay	9.6+/-6.5	11.4+/-6.2	0.709	10+/-6.3	10.2+/-6.6	0.885

TABLE III: ASSOCIATION OF DEMOGRAPHIC AND MEDICAL PROFILE WITH OUTCOMES OF COVID-19 PATIENTS ADMITTED IN ICU

Associated factors	Outcome		P-values	
	Expired	Discharge		
Age groups (in years)	Less than 40	0(0%)	5(7.8%)	0.009 ^x
	41-60	9(23.7%)	29(45.3%)	
	More than 60	29(76.3%)	30(46.9%)	
Gender	Female	16(42.1%)	28(43.8%)	0.871 ^x
	Male	22(57.9%)	36(56.2%)	
Diabetes mellitus	23(60.5%)	34(53.1%)	0.467 ^x	
Hypertension	27(71.1%)	31(48.4%)	0.026 ^x	
Asthma	2(5.3%)	3(4.7%)	0.896 ^x	
COPD	2(5.3%)	2(3.1%)	0.591 ^F	
IHD	18(47.4%)	10(15.6%)	0.001 ^x	

patient outcomes and demographic/ medical profile. Age showed a significant effect on mortality in these patients. The mortality rate was higher in older patients (P-value: 0.009). There was a slightly higher incidence of mortality among males when compared to females. Mortality was seen more among patients have with co-morbidities, especially patients with hypertension, DM, and IHD. Comparison of electrolytes levels between expired and survived patients was also stated. (Table III)

Comparisons of Electrolyte levels

Sodium	134.8±5.96	134.9±5.64	0.916 ^T
Potassium	4.26±0.74	4.08±0.69	0.223 ^T
Chloride	101.66±6.93	102.05±6.31	0.773 ^T
Magnesium	2.18±0.43	2.07±0.42	0.190 ^T
Calcium	8.21±0.70	8.30±0.71	0.551 ^T
Phosphorus	4.37±2.78	3.83±1.95	0.256 ^T

Chi-square test= χ ; Fisher exact test=F; Independent t-test=t; significance level<0.05

DISCUSSION

The present study aimed to report the pattern of electrolytes abnormalities. Further to evaluate the impact of these abnormalities in terms of in-hospital outcomes on ICU admitted COVID-19 patients. There is a need to study electrolytes imbalances and their management in a current pandemic. So far, a few studies have been reported electrolyte abnormalities in COVID-19 patients. Studies conducted by Sultana R et al.², Sarvazadet H 2020⁵, and Tezcan ME 2020³ assessed electrolyte imbalances in patients with COVID-19. These studies revealed that COVID-19 is strongly associated with low levels of sodium, potassium, calcium, magnesium, etc. From these studies, it was discovered that 38 to 77%^{2,3,5} patients experienced hyponatremia, 7% to 50%^{2,3,5} experienced hypokalemia, 9% to 28%^{2,3} experienced hypocalcemia and 15% to 32%^{2,5} experienced hypomagnesemia. These results are comparable with our study findings and some other studies^{6,11}. Previous studies were limited to sodium and potassium levels. Interestingly, we have reported electrolytes abnormalities in patients admitted in ICU with severe Covid-19 also evaluated the impact of these abnormalities on patients' in-hospital outcomes and reported previously neglected parameters such as hypermagnesemia, making our study unique novel.

The findings of our study showed that males had a more significant proportion in the cases with COVID-19 compared to females. The results concur with the investigations performed in neighboring countries, i.e., Chinese and Bangladeshi. They also revealed a strong link between hypertension and diabetes with a higher rate of morbidity, severity, and poor prognosis in COVID patients^{2,18,19}. Regarding high sodium levels, this study revealed that none of the patients had experienced hypernatremia which is in disagreement with previous studies^{5,7,15}. The studies showed that high sodium levels led to a more extended period of intensive care and increased mortality risk. With regards to low levels of sodium, 45.1% of patients had experienced hyponatremia in the present study cohort. The other two studies had reported similar findings for low sodium and severity of COVID-19^{6,20}. Considering potassium levels, a study showed that hypokalemia is also challenging to manage because of the deprivation of the angiotensin-converting enzyme.²¹ The pathophysiology of SARS-CoV-2 reflects that decreased angiotensin-2 increases faecal potassium levels leading to hypokalemia.⁵ In our study, only 17.6% of patients with severe COVID-19 experienced hypokalemia. Whereas Chen D et al.¹¹ revealed a higher frequency of hypokalemia in these patients. Therefore, it is advised that continuous monitoring and substitution of potassium levels might help reduce the risk of imbalance in electrolytes and their related complications.

Magnesium blocks calcium induction into immune-

competent cells. Therefore, it is known as a calcium channel blocker. It inhibits systemic inflammation by inhibiting the production of cytokines and interleukins. Hence, hypomagnesemia exacerbates inflammation in COVID-19.⁵

Iotti S 2020²² discovered stress as one of the causes of hypomagnesemia in a current pandemic. As with stress, catecholamines and corticosteroids push magnesium to extracellular space, increasing faecal magnesium. In this study, 12.7% of patients experienced hypomagnesemia, and 40.2% of patients experienced hypermagnesemia. Duan J et al. establish that potassium, sodium, and chloride levels can be predictors for severe COVID-19 disease¹².

The prognosis factors such as hospitalization and ICU admissions were also investigated in previous studies. It was revealed that high CRP levels, old age, and low white blood cells were associated with hospitalization²³, whereas prolonged hospitalization was linked with high ferritin levels and LDH²⁴. Moreover, a systematic review found that hyponatremia and hypokalemia are related to severe COVID⁶, whereas high sodium, chloride, and potassium were linked to progression towards severe COVID¹². Our study found that low baseline sodium, chloride, and calcium levels were the significant predictors of higher incidence of mortality, higher ICU admission, prolonged stay and Mechanical ventilation requirement, and more extended hospital stays. The studies above showed that electrolyte abnormality is strongly associated with severe COVID-19 and poor prognosis. However, sodium levels are commonly found low in the studies described above. Hence, targeting and monitoring sodium levels could help better clinical outcomes for the patients and be less challenging for physicians. It is also important to assess other electrolyte abnormalities that impact clinical outcomes, such as Hypocalcemia, hypermagnesemia, and hyperchloremia. The present study emphasizes the monitoring of these electrolytes for better treatment outcomes and preventing complication of the disease.

The current study has a limited sample size. Based on the findings of this research, it is recommended to explore in-depth information on a larger sample size for the longer follow-up to achieve generalizability. Moreover, causes and impact of electrolytes abnormality should be established, and a wide variety of electrolytes should be assessed for a better prognosis of critically ill Covid-19 patients.

CONCLUSION

The study showed that most patients with severe Covid-19 had electrolyte abnormalities at admission into the intensive care unit. Abnormalities were found in 90% of the ICU admitted Covid-19 patients, while 70% had more than two electrolyte imbalances. Hyponatremia, hypermagnesemia, and Hypocalcemia

were the most common electrolytic abnormalities encountered in this study. The electrolyte imbalances are strongly linked with poor clinical outcomes and factors such as life support requirement, prolonged ICU stays, and increased risk of mortality.

Ethical permission: Centre of International Research Sciences Karachi, ERC letter No. ERC-CIRS-2021-07, Dated: 02-04-2021.

Conflict of Interest: There is no conflict of interest among the authors.

Financial Disclosure / Grant Approval: There was no funding agency.

Data Sharing Statement: The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions

AUTHOR CONTRIBUTIONS

All authors have contributed equally: Haroon A, Abbas SA, Khan A, Ali M, Qazi R, Kumar A.

A. Substantial contribution to conception or design of the work; or the acquisition, analysis, or interpretation of data for the work

B. Drafting of the work or revising it critically for important intellectual content

C. Final approval of the version to be published

D. Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

REFERENCES

1. Omar S, Baker D, Siebert R, Joubert I, Levy B, Paruk F et al. The role of laboratory testing in hospitalised and critically ill COVID-19-positive patients. *South Afr J Crit Care*. 2020; 36(1): 14-7.
2. Sultana R, Ahsan AA, Fatema K, Ahmed F, Saha DK, Saha M et al. Pattern of electrolytes in a cohort of critically ill COVID-19 patients. *BIRDEM Med J*. 2020; 10: 46-50. doi: 10.3329/birdem.v10i0.50980.
3. Tezcan ME, Gokce GD, Sen N, Kaymak NZ, Ozer R. Baseline electrolyte abnormalities would be related to poor prognosis in hospitalized coronavirus disease 2019 patients. *New Microbes New Infect*. 2020; 37: 100753. doi: 10.1016/j.nmni.2020.100753.
4. Gupta L, Jalang'o GA, Gupta P. Nutritional management and support in COVID-19: Emerging nutrigenetics. *J Pak Med Assoc*. 2020; 70(suppl 3): S124-S130. doi: 10.5455/JPMA.31.
5. Sarvazad H, Cahngaripour S, Roozbahani NE, Izadi B. Evaluation of electrolyte status of sodium, potassium and magnesium, and fasting blood sugar at the initial admission of individuals with COVID-19 without underlying disease in Golestan Hospital, Kermanshah. *New Microbes New Infect*. 2020; 38: 100807. doi: 10.1016/j.nmni.2020.100807.
6. Lippi G, South AM, Henry BM. Electrolyte imbalances in patients with severe coronavirus disease 2019 (COVID-19). *Ann Clin Biochem*. 2020; 57(3): 262-5. doi: 10.1177/0004563220922255.
7. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX et al. Clinical characteristics of coronavirus disease 2019 in China. *New Engl J Med*. 2020; 382(18): 1708-20. doi: 10.1056/NEJMoa2002032
8. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020; 395(10223): 497-506. doi: 10.1016/S0140-6736(20)30183-5.
9. Nair V, Niederman MS, Masani N, Fishbane S. Hyponatremia in community-acquired pneumonia. *Am J Nephrol*. 2007; 27(2): 184-90. doi: 10.1159/000100866.
10. Wang S, Ma P, Zhang S, Song S, Wang Z, Ma Y et al. Fasting blood glucose at admission is an independent predictor for 28-day mortality in patients with COVID-19 without previous diagnosis of diabetes: a multi-centre retrospective study. *Diabetologia*. 2020; 63(10): 2102-11. doi: 10.1007/s00125-020-05209-1.
11. Chen D, Li X, Song Q, Hu C, Su F, Dai J et al. Assessment of hypokalemia and clinical characteristics in patients with coronavirus disease 2019 in Wenzhou, China. *JAMA Netw Open*. 2020; 3(6): e2011122. doi: 10.1001/jamanetworkopen.2020.11122.
12. Duan J, Wang X, Chi J, Chen H, Bai L, Hu Q et al. Correlation between the variables collected at admission and progression to severe cases during hospitalization among patients with COVID-19 in Chongqing. *J Med Virol*. 2020; 92(11): 2616-22. doi: 10.1002/jmv.26082.
13. Li X, Hu C, Su F, Dai J. Hypokalemia and clinical implications in patients with coronavirus disease 2019 (COVID-19). (Preprint February 29, 2020) *MedRxiv*. 2020. doi: <https://doi.org/10.1101/2020.02.27.20028530>
14. Liu Y, Yang Y, Zhang C, Huang F, Wang F, Yuan J et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci*. 2020; 63(3): 364-74. doi: 10.1007/s11427-020-1643-8.
15. Sjöström A, Rysz S, Sjöström H, Höybye C. Hyponatremia is common in patients with severe COVID-19 and indicates a poor prognosis. *Research Square*. Preprint. 2020.
16. Garcia PDW, Fumeaux T, Guerci P, Heuberger DM, Montomoli J, Roche-Campo F et al. Prognostic factors associated with mortality risk and disease progression in 639 critically ill patients with COVID-19 in Europe: Initial report of

- the international RISC-19-ICU prospective observational cohort. *E Clin Med.* 2020; 25: 100449.
17. Organization WH. Clinical management of COVID-19: interim guidance, 27 May 2020. World Health Organization, 2020.
 18. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX et al. Clinical characteristics of 2019 novel coronavirus infection in China. *MedRxiv.* 2020.
 19. Hu Y, Sun J, Dai Z, Deng H, Li X, Huang Q et al. Prevalence and severity of coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis. *J Clin Virol.* 2020; 104371. doi: 10.1016/j.jcv.2020.104371.
 20. Zhang W, Lu S, Zhang M, Zheng H, Huang Y, Chen S et al. Correlation between hyponatremia and the severity of coronavirus disease 2019. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue.* 2020; 32 (7): 774-8. doi: 10.3760/cma.j.cn121430-20200331-00417.
 21. Mabillard H, Sayer JA. Electrolyte Disturbances in SARS-CoV-2 Infection. *F1000Research.* 2020; 9: 587. doi: 10.12688/f1000research.24441.2.
 22. Iotti S, Wolf F, Mazur A, Maier JA. The COVID-19 pandemic: is there a role for magnesium? Hypotheses and perspectives. *Magnes Res.* 2020; 33(2): 21-7. doi: 10.1684/mrh.2020.0465.
 23. Hou W, Zhang W, Jin R, Liang L, Xu B, Hu Z. Risk factors for disease progression in hospitalized patients with COVID-19: a retrospective cohort study. *Infect Dis(Lond).* 2020; 52(7): 1-8. doi: 10.1080/23744235.2020.1759817.
 24. Tezcan ME, Doğan Gökçe G, Ozer RS. Laboratory abnormalities related to prolonged hospitalization in COVID-19. *Infect Dis(Lond).* 2020; 52(9): 666-8. doi:10.1080/23744235.2020.1776381.



AUTHOR AFFILIATION:

Dr. Anita Haroon (*Corresponding Author*)

Assistant Professor
Baqai Medical University
OMI Hospital Karachi, Sindh-Pakistan.
Email: dr.anitaharoon@gmail.com

Dr. Syed Ali Abbas

Department of Pulmonology and Critical Care
Dr. Ziauddin University Hospital
Karachi, Sindh-Pakistan.

Dr. Amanullah Khan

Assistant Professor of Medicine
Baqai Medical University Karachi, Sindh-Pakistan.

Dr. Momina Ali

House officer
Dr. Ziauddin University Hospital
Karachi, Sindh-Pakistan.

Dr. Rija Qazi

House officer
Dr. Ziauddin University Hospital
Karachi, Sindh-Pakistan.

Dr. Ajeet Kumar

Professor of Medicine
Jinnah Medical College Hospital
Karachi, Sindh-Pakistan.