

Assessment of Thyroid Profile in Diabetes Mellitus Type II Patients at LUMHS, Jamshoro/ Hyderabad

Kavita Bai, Abdullah Abbassi, Ramesh Kumar Suthar, Rasheeda Qasmi,
Urooj Bhatti, Keenjhar Rani

ABSTRACT

OBJECTIVES: To assess status of thyroid functions among the type 2 diabetic patients.

METHODOLOGY: This cross-sectional comparative study conducted in Department of Physiology LUMHS, in collaboration with Diabetic clinic and diagnostic research laboratory of LUMHS, Jamshoro from September 2015 to February 2016 with non-probability sampling technique. The volunteers recruited in the study were one hundred two (n, 102) type 2 diabetic patients and one hundred two (n, 102) non diabetic individuals. Thyroid stimulating hormone (TSH) and thyroid hormones (T3 and T4) were measured using autoanalyzer by Cobas e411, Roche. The data analyzed on SPSS version 16.0 and the p value less than 0.05 were reflected statistically significant.

RESULTS: Mean age of study participants was 45.85 years \pm 9.08 in this research study. The mean and standard deviation of T₃ among type 2 diabetic and non- diabetic groups were 1.144ng/ml \pm 0.38 and 1.145ng/ml \pm 0.36 respectively with insignificant difference (p value = 0.98). However, the mean \pm standard deviation of T4 levels among type 2 diabetic and non- diabetic groups were 7.13 μ g/dl \pm 1.95 and 7.33 μ g/dl \pm 2.0 respectively with non-significant difference (p value = 0.98). Instead, there was statistically noteworthy difference of TSH between type 2 diabetic and non-diabetic people (p value revealed to be <0.01). Mean \pm standard deviation of TSH was 6.42 μ U/ml \pm 1.36 and 1.87 μ U/ml \pm 1.30 among type 2 diabetic patients and non- diabetic healthy individuals respectively. Thyroid dysfunctions found to be 18% among type 2 DM patients.

CONCLUSION: It has been concluded from this research that there is a significant variance of TSH levels among type 2 diabetic patients and non-diabetic healthy individuals.

KEY WORDS: Type 2 diabetes mellitus, Non-diabetic healthy individuals, thyroid functions

This article may be cited as: Bai K, Abbassi A, Suthar RK, Qasmi R, Bhatti U, Rani K. Assessment of Thyroid Profile in Diabetes Mellitus Type 2 Patients at LUMHS, Jamshoro/ Hyderabad. J Liaquat Uni Med Health Sci. 2021;20(02):109-12. doi: 10.22442/jlumhs.2021.00721

INTRODUCTION

Type 2 diabetes mellitus (T2DM) is the universal rampant condition that is linked to peaked health expenses, as well as curtailed quality of life¹. Diabetes mellitus (DM) is a pool of metabolic conditions, characterized by hyperglycemia that result from either due to the defects in secretion of insulin or resistance of insulin action or may be both. The wide-reaching prevalence of DM is estimated to be 366 million people worldwide².

Thyroid dysfunctions are more expected to arise among DM patients than general healthy population. It is proposed by the research that assessment of thyroid functions should be done in high-risk conditions as a slice of wide-ranging management among the T2DM patients³.

The ratio of disturbed thyroid functions is greatly extensive among diabetic population and estimated to be between 6.9% and 16 percent DM has impact on thyroid function on two locations; first at the level of hypothalamic regulation of release of thyroid stimulating hormone (TSH) and other at the level of peripheral tissue where conversion of thyroxine (T4) to

tri-iodothyronine (T3) take place. Both DM and hypothyroidism are major endocrine disorder leading to hyperlipidemia,⁴ so T2DM remains associated to augmented cardiac morbidity and ultimately to increased mortality rate⁵. Both DM and thyroid disorders are the two utmost combined syndromes of endocrine system. Great proportion of T2DM patients suffer from never diagnosed thyroid illnesses. Subclinical hypothyroidism has been discovered as the furthestmost common thyroid ailment among type 2 diabetic patients⁶.

Dyslipidemia, elevated LDL levels, and osteoporosis are most frequent among patients with disturbed thyroid function when compared with those having normal thyroid functions. Thyroid dysfunctions remain underdiagnosed among elderly type 2 diabetic patients⁷. The metabolic control of diabetic patients may be impaired among untreated cases and this association can have important repercussions on the outcome of both of these disorders⁸. This is typically challenging to diagnose clinically thyroid deviations among the diabetic individuals because features of thyroid deviations may pretend symptoms of DM and its complications. Consequently, assessing

prevalence of thyroid dysfunctions among the patients of T2DM would support improved control of DM as well as its complications⁹. Thyroid function tests should be recommended for screening yearly in DM patients to be distinguished from subclinical asymptomatic thyroid disorders. The rationale of this research project was to generate evidence about eminence of thyroid profile among T2DM patients and to produce cognizance among such people regarding significance of screening thyroid functions among T2DM patients. In upcoming times, this research study may benefit to improve the strategies as well as policies to screen such diabetic cases earlier and to be treated and coped timely.

METHODOLOGY

This cross-sectional comparative study conducted in Department of Physiology at LUMHS, Jamshoro in collaboration with Diabetic OPD of Liaquat University Hospital Hyderabad / Jamshoro and Diagnostic and Research Laboratory LUMHS, Jamshoro from September 2015 to February 2016. Sampling technique was non probability sampling. According to a research published in JPMA¹⁰, prevalence of type 2 DM patients in Pakistan was 11.77%, taking 95% confidential interval and 5% margin of error, calculated sample size was 102 (n=102) type 2 DM patients and for comparison 102 non diabetic included. So that, total 204 volunteers were included in this research project and separated into two sets, (102 type 2 diabetic) and (102 non diabetic). Both male and female, young adults diagnosed patients of T2DM who never taken the drugs affecting thyroid profile were recruited for this research project. Inclusion Criteria for controls were both male and female healthy young individuals having blood sugar fasting (BSF) normal after two readings of BSF on two different occasions according to WHO criteria. The persons suffering from former endocrine pathology, diagnosed cases of type 1 DM, those who are taking lipid dropping drugs, having pregnancy, with past history of thyroid dysfunctions, taking the drugs altering thyroid functions as well as, also the diseases disturbing levels of thyroid hormones i.e., Cerebrovascular, cardiovascular, neurological and hepatic diseases, chronic kidney disease, asthmatics were excluded for this research study.

After subsequent consents and later taking aseptic measures; five ml intravenous blood samples were taken from the volunteers meeting the inclusion criteria. Then for biochemical analysis purpose, these blood samples sent to the laboratory to measure the TSH, T3 and T4 levels. Serum parted and then deposited at temperature of -20⁰c till investigated. Biochemical analyzer used for biochemical analysis was Cobas e411 by Roche, by means of electro chemiluminescence skill rendering to the instructions of producers. Patients were labeled as subclinical

hypothyroidism when T3 levels were 0.58 to 1.59 ng/ml, T4 levels found 4.87^o 11.27 µg/dl with TSH > 4.00 µU/ml. The patients considered as suffering from clinical hypothyroidism if T3 levels revealed less than 0.58 ng /ml, T4 levels less than 4.87 µg/dl and TSH more than 4.0 µU/ml.

Data analyzed on statistical package of social sciences (SPSS) version 16.0. Frequencies with percentage calculated for descriptive data while quantitative variables documented as mean with standard deviation. Outcome modifiers organized by stratifying both age as well as gender. For relating qualitative data, chi square test applied. The p-value ≤ 0.05 were reflected statistically significant.

RESULTS

Entire 204 (n=204) volunteers were included in this research project and separated into two sets, (102 type 2 diabetic) and (102 controls). In each set, further males and females separated equally. Mean of the age with standard deviation of study participants was 45.85years ± 9.08. The mean ± standard deviation of T₃ among type 2 diabetic and control non diabetic groups were found to be 1.14ng/ml ± 0.38 and 1.14ng/ml±0.36 respectively with unimportant variance (p value = 0.98). However, the mean of T4 levels among type 2 diabetic and non-diabetic groups were 7.31µg/dl±1.95 and 7.35±2.0 respectively with non-significant alteration (p value = 0.98). Instead, it has been observed that statistically significant variance exists of TSH between type 2 DM group and non-diabetic controls with p value less than 0.01with the mean values 6.42µU/ml±1.36 and 1.87µU/ml±1.30 among type 2 DM patients and control groups correspondingly **Table I**.

In this study, occurrence of thyroid disease revealed to be 18% among 102 type 2 diabetics while maximum among female gender (11.8%). Among the diabetics, 7.8% were suffering from subclinical hypothyroidism. Two percent (2%) diagnosed as subclinical hyperthyroidism as well as one percent as hyperthyroidism. Female diabetics are more disposed to develop thyroid dysfunctions. **Table II**.

TABLE I: MEAN AGE, T3, T4 AND TSH AMONG THE STUDY POPULATION (n=204)

Study Population	N	Parameters	Mean± Std Deviation	p-value
Type 2 DM Patients	102	Age (in years)	46.49±9.92	0.317
Non-diabetic	102		45.22±8.15	
Type 2 DM Patients	102	T3(ng/ml)	1.144±0.38	0.98
Non-diabetic	102		1.145±0.34	
Type 2 DM Patients	102	T4(µg/dl)	7.31±1.95	0.89
Non-diabetic	102		7.35±2.05	
Type 2 DM Patients	102	TSH(µU/ml)	6.42±1.36	0.001
Non-diabetic	102		1.87±1.30	

Table II: THYROID STATUS AMONG TYPE 2 DIABETIC GROUP (n = 102)

Thyroid status	Type 2 DM patients			P value
	Male n = 51	Female n=51	Total n=102	
Euthyroidism	48(94.1%)	42(82.4%)	90(88.2%)	0.029*
Subclinical hypothyroidism	2(3.9%)	6(11.8%)	8(7.8%)	
Hypothyroidism	1(2.0%)	2(3.9%)	3(2.9%)	
Hyperthyroidism	0	1(2.0%)	1(1.0%)	

DISCUSSION

DM and thyroidal illness both are the major global issues related to endocrine system apparent in largely population. The association between DM and thyroid dysfunction had been documented and published previously. Screening for thyroid profile among DM patients can possibly improve incident recognition as well as timely management, ultimately foremost towards better thyroid specific management as well as prevention from further DM complications¹¹.

Among general population, thyroid illnesses range from 6.6 to 13.4 percent.⁵ Instead, among DM patients, the prevalence of thyroid disorders observed to be varying from 10 to 24%. However, this study found the occurrence of entirely types of undiagnosed thyroid disorders to be 18.3%⁶. Peak level of TSH with low levels of T3 has been detected in T2DM. Routine monitoring of thyroid in DM patients is necessary, and management of thyroid dysfunction may be a potential therapeutic strategy of DM¹².

In this study, the prevalence of subclinical hypothyroidism is more in T2DM people. This is in association to the study of Elgazar EH 2019⁹ and Alsolami AA et al¹³ who concluded greater prevalence of thyroid troubles among patients suffering from T2 DM. And this dysfunction has been revealed as, in direct proportion with HbA1c levels. This might propose that the deprived glycemic mechanism can possibly have a part in the expansion to thyroid dysfunctions in the patients suffering from T2DM. Subclinical hypothyroidism has been found to be utmost prevalent kind of thyroid disorders in DM patients⁹. Hypothyroidism is a recognized risk feature for developing coronary artery disease (CAD) and is observed to be related with augmented all-cause mortality independent of CAD risk elements. T2DM patients were commonly seen to be at augmented risk of emerging hypothyroidism. Adequate management and control of T2DM might lessen the risk of emerging hypothyroidism¹³. According to a research study, prevalence of subclinical hypothyroidism and overt hypothyroidism was 23/100 (23%) and 3/100 (3%), respectively¹⁴.

Tiwari HK 2019¹⁵ also revealed that Subclinical hypothyroidism and overt hypothyroidism was the most common thyroid abnormality in Type 2 DM.

Thyroid dysfunction was associated with worsening dyslipidemia in T2DM. The mean body mass index was high in diabetic patients with thyroid dysfunction. The hemoglobin A1c levels in patients who had thyroid dysfunction were high. Different research studies revealed that hypothyroidism, particularly subclinical variety, is seen in type 2 diabetic patients. Therefore, routine screening is advised^{16,18}, but one study conducted at South India revealed that thyroid function profile is not altered among the T2DM patients when compared to nondiabetics¹⁷.

Conferring to this research study, the occurrence of subclinical hypothyroidism observed to be advanced and prevalent among female T2 DM patients in comparison with general apparently healthy population. This innovation is consistent with the research by different authors i.e., Lal B 2016¹⁸ and Mishra S 2016¹⁹ revealed greater pervasiveness of thyroid dysfunctions among females having DM. Among hypothyroid individuals, 23.08 percent and 10 percent were male and females respectively, with noteworthy variance (p=0.017), however among these hypothyroid people, 4 %observed to be suffering from clinical hypothyroidism while 9 % cases of subclinical hypothyroidism detected¹⁸. It is beneficial to screen T2DM patients for thyroid issues and this could make progress in early case detection as well as timely management, indirectly leading to improved thyroid related outcomes with the prevention of the DM complications.

CONCLUSION

It has been concluded from this research study that T2DM is one of the risk factors for thyroid dysfunctions and guidelines should be made to diagnose and manage such co morbid cases timely.

Ethical permission: Liaquat University of Medical & Health Sciences ERC letter No. LUMHS/REC/=374, dated 13-08-2015.

Conflict of interest: There is no conflict of interest.

Funding: There was no any funding agency.

AUTHOR CONTRIBUTIONS

Bai K: Concept, data collection & statistics
 Abbassi A: Critical revision & suggestion
 Suthar RK: Data collection & literature search
 Qasmi R: Critical revision & suggestion
 Bhatti U: Data analysis
 Rani K: Data collection

REFERENCES

- Bellou V, Belbasis L, Tzoulaki I, Evangelou E. Risk factors for type 2 diabetes mellitus: An exposure-wide umbrella review of meta-analyses. *PLoS ONE*. 2018; 13(3): e0194127. doi:10.1371/journal.pone.0194127.
- Sunkari S, Godbole G, Ashok P, Kharche J, Joshi A. Association of hypothyroidism and type 2 diabetes

- in patients attending tertiary care hospital of Pune City. *Natl J Physiol Pharm Pharmacol*. 2019; 9(4): 328-30. doi:10.5455/njppp.2019.9.1236811022019.
3. Subekti I, Pramono LA, Dewiasty E, Harbuwono DS. Thyroid dysfunction in type 2 diabetes mellitus patients. *Acta Med Indones*. 2017; 49(4): 314-323.
 4. Chand G, Garg P, Oberoi KK, Sandhu S. Study of dyslipidemia in type 2 diabetes mellitus patients with and without associated hypothyroidism. *Int J Sci Res*. 2019; 8(3): 66-68.
 5. Zweck E, Scheiber D, Jelenik T. Insulin Resistance and Type 2 Diabetes Mellitus Are Associated with Impaired Mitochondrial Function in Human Ventricular Myocardium. *Diabetes*. 2018;67(1): 284. doi:10.2337/db18-284.
 6. Khan RN, Ullah R, Ahmad I, Ghaffar T. Frequency of undiagnosed thyroid disorders in patients with type 2 diabetes mellitus at a tertiary care center. *J Postgrad Med Inst*. 2018; 32(2): 162-165.
 7. Zhu Y, Xu F, Shen J, Liu Y, Bi C, Liu J, et al. Prevalence of thyroid dysfunction in older Chinese patients with type 2 diabetes-A multicenter cross-sectional observational study across China. *PLoS One*. 2019; 14(5): e0216151. doi: 10.1371/journal.pone.0216151.
 8. Biondi B, Kahaly GJ, Robertson RP. Thyroid Dysfunction and Diabetes Mellitus: Two Closely Associated Disorders. *Endocr Rev*. 2019; 40(3): 789-824. doi: 10.1210/er.2018-00163.
 9. Elgazar EH, Esheba NE, Shalaby SA, Mohamed WF. Thyroid dysfunction prevalence and relation to glycemic control in patients with type 2 diabetes mellitus. *Diabetes Metab Syndr*. 2019; 13(4): 2513-2517. doi:10.1016/j.dsx.2019.07.020.
 10. Meo SA, Zia I, Bukhari IA, Araisn SA. Type 2 diabetes mellitus in Pakistan: Current prevalence and future forecast. *J Pak Med Assoc*. 2016; 66(12): 1637-42.
 11. Hussain T, Barik BS, Nayak AR, Das S, Khadanga UK, Yadav VS, et al. Prevalence and predictors of thyroid dysfunction among patients with type 2 diabetes mellitus attending a tertiary care hospital in an urban area of Bhubaneswar, Odisha. *Thyroid Res Pract*. 2019; 16(1): 26-35.
 12. Zhao W, Li X, Liu X, Lu L, Gao Z. Thyroid Function in Patients with Type 2 Diabetes Mellitus and Diabetic Nephropathy: A Single Center Study. *J Thyroid Res*. 2018; Article ID: 9507028. doi:10.1155/2018/9507028.
 13. Alsolami AA, Alshali KZ, Albeshri MA, Alhassan SH, Qazli AM, Almalki AS, et al. Association between type 2 diabetes mellitus and hypothyroidism: a case-control study. *Int J Gen Med*. 2018; 11: 457-461. doi: 10.2147/IJGM.S179205.
 14. Pramanik S, Ghosh S, Mukhopadhyay P, BhattacharjeeR, MukherjeeB, Mondal SA, et al. Thyroid Status in Patients with Type 2 Diabetes Attending a Tertiary Care Hospital in Eastern India. *Indian J Endocrinol Metab*. 2018; 22(1): 112-115. doi:10.4103/ijem.IJEM_572_17.
 15. Tiwari HK, Kumar P. A case control study to evaluate the prevalence of thyroid dysfunction in Type-2 Diabetes Mellitus. *J Adv Med Dent Sci Res*. 2019; 7(6): 57-9.
 16. Sunkari S, Godbole G, Ashok P, Kharche J, Joshi A. Association of hypothyroidism and type 2 diabetes in patients attending tertiary care hospital of Pune City. *Natl J Physiol Pharm Pharmacol*. 2019; 9(4): 328-330. doi:10.5455/njppp.2019.9.1236811022019.
 17. Menon UV, Bhavani N, Jayakumar RV, Kumar H, Nair V, Praveen VP, et al. Prevalence of thyroid disorders is not different in type 2 diabetes mellitus compared to nondiabetics in South India. *Int J Diabetes Dev Ctries*. 2019; 39(2): 262-7. doi:10.1007/s13410-018-0672-8
 18. Lal B, Pandey M, Rao S, Mathur S, Rishi JP. Prevalence of Thyroid Disorders in Type 2 Diabetes Mellitus. *Ann Int Med Dent Res*. 2016; 2(1): 216-9.
 19. Mishra S, Mishra S, Behera BK. Thyroid dysfunction in patients with Type-2 diabetes mellitus in Eastern India. *J Pharm Biomed Sci*. 2016; 06(05): 311-314.

AUTHOR AFFILIATION:

Dr. Kavita Bai (*Corresponding Author*)
Assistant Professor, Department of Physiology
Indus Medical College
Tando Muhammad Khan, Sindh-Pakistan.
Email: baidrkavita@gmail.com

Dr. Abdullah Abbassi
Professor, Department of Physiology
Muhammad Medical College
Mirpurkhas, Sindh-Pakistan.

Dr. Ramesh Kumar Suthar
Assistant Professor, Department of Medicine
Indus Medical College

Tando Muhammad Khan, Sindh-Pakistan.

Dr. Rasheeda Qasmi
Professor, Department of Physiology
Muhammad Medical College
Mirpurkhas, Sindh-Pakistan.

Dr. Urooj Bhatti
Lecture, Department of Physiology
Liaquat University of Medical and Health Sciences
(LUMHS), Jamshoro, Sindh-Pakistan.

Dr. Keenjhar Rani
Assistant Professor, Department of Physiology
LUMHS, Jamshoro, Sindh-Pakistan.