

Knowledge, Attitude and Practices of Dengue Fever among Rural Population of Gujrat

Syed Muhammad Ali Shah, Kanza Malik, Ali Malik, Maryam Abid, Sofia Bano

ABSTRACT

OBJECTIVE: To acquire and collect information regarding Dengue Fever (DF) awareness in rural population.

METHODOOGY: A descriptive cross sectional observational study was conducted among 100 individuals from rural population of Gujrat from March to May 2017 and non-probability cluster sampling was done. All consenting individuals aged above 18 years and above encountered in selected rural areas interviewed using a structured, pre-tested questionnaire. Data regarding age, gender, occupation, mode of spread, vector, prevention, management and outcomes was collected. Data was analyzed using SPSS version 20.0. Results are expressed as mean±SD for continuous variables such as age and number (percentage) for categorical data such as gender.

RESULTS: Mean age was 30.24±16.57 years, 63% male and 37% female. Most of respondents did outdoor jobs and were young. Respondents had good knowledge regarding vector, its breeding places, symptoms and outcome of disease but lacked knowledge in breeding season, treatment options, management and prevention of dengue fever.

CONCLUSION: The rural population of Gujrat has good knowledge regarding mode of spread, symptoms of dengue fever and where to get treatment however they lacked knowledge in management, outcome and prevention of dengue fever.

KEY WORDS: Dengue Fever, Knowledge, Prevention, Rural

This article may be cited as: Shah SMA, Malik K, Malik A, Abid M, Bano S. Knowledge, Attitude and Practices of Dengue Fever among Rural Population of Gujrat. J Liaquat Uni Med Health Sci. 2018;17(03):185-9. doi: 10.22442/jlumhs.181730575

INTRODUCTION

Dengue Fever (DF) is an important public health problem in Pakistan and its incidence has grown dramatically around the world claiming thousands of lives. Its first outbreak was reported in Pakistan in 1994 and an epidemic has been declared since 2010 and thousands of patients have been reported with hundreds of deaths attributable to this disease¹. The incidence of dengue fever is 0.5 per 100,000 in year 2016 as reported by Government of Pakistan². There is a transition in pattern of disease from mild to severe and multiple factors have been reported including lack of resources, climate change, natural disasters and resistance of mosquitoes to insecticides.³ The main causes and barriers regarding dengue control depicted in another study in Pakistan included congested streets and houses, large population, presence of mosquito breeding sites, and a passive role of government regarding community awareness regarding DF⁴.

Multiple knowledge, attitude and practices (KAP) studies have been conducted in Pakistan regarding dengue fever. One study conducted in Karachi

demonstrated that the knowledge regarding dengue fever is inadequate among people with low socioeconomic group with only 35% peoples having adequate knowledge⁵. In another KAP study conducted among teachers concluded that they have adequate knowledge regarding DF⁶. In another study conducted among university students of interior Sindh Bota R 2014⁸ demonstrated that almost half of the participants could not identify the Aedes mosquito as transmission source, appearance of mosquito, breeding place and bite time⁷. In another study in D.I. Khan conducted among patients of DF and their attendants showed that they have a very low knowledge regarding DF⁸. Another study in Karachi concluded that knowledge of DF, threat perception, and self-efficacy are significant predictors of prevention of DF among local population⁹. About 67.5% of population of Pakistan lives in rural areas¹⁰ and Pakistan had literacy rate of 56% in 2014¹¹ which is very low. Most of people in rural areas have poor life style and economic condition, less access to medical services along with less literacy rate which are important factors in decreased community awareness regarding DF and leading to

negligible preventive measures for spread of DF. This is a point of concern as it may lead to increased mortality in rural population and there is no documented evidence regarding awareness and practices of dengue fever among rural population. Thus this study was aimed at accessing the frequency of knowledge of rural population regarding mode of spread, prevention and outcome of DF.

METHODOLOGY

A descriptive cross sectional observational study was conducted among 100 individuals from rural population of district Gujrat. Different rural areas of Gujrat were visited by a team of medical students and doctors for the purpose of data collection and probability cluster sampling was done from selected communities and localities visited for the survey which was dependent on a number of factors such as convenience of commute and distance from Aziz Bhatti Shaheed Teaching Hospital Gujrat. Study was conducted from 1st March 2017 to 31st May 2017.

Knowledge attitude and practices among rural population are assessed by our questionnaire. All consenting individuals aged above 18 years and above encountered in selected rural areas were interviewed using a structured, pre-tested questionnaire. The questionnaire was thoroughly discussed before hand to minimize interview bias. Informed consent was taken from all the respondents and confidentiality was ensured throughout the study. The questionnaire and consent forms were translated into Urdu for the ease of respondents. Every individual was given a total score based on this variable in order to categories them into age, gender, education, occupation, mode of spread, vector, prevention, management and outcomes.

Any scoring with higher percentage gave us information about the knowledge, attitude and practices of dengue fever among rural population of Gujrat. We have to choose a number of variables. This strategy was used because a single variable cannot accurately illustrate an individual's knowledge attitude and practices of dengue fever.

Data was analyzed using SPSS version 20.0. Results are expressed as mean±SD for continuous variables such as age and number (percentage) for categorical data such as gender.

RESULTS

A total of 100 individuals were selected to participate in this survey from different localities in rural suburbs of district Gujrat. The response rate was 100% and all respondents were cooperative and successfully interviewed. Mean age of respondents was

30.24±16.57 years. Percentage of respondents according to age group; 16-30 years in 68%, 30-45 years in 16%, 40-60 years in 9% and 61-75 years in 7% of respondents. Majority respondents were male; 63% of total while female respondents were 37%.

Respondents belonged to different occupations with top of list students (36%) followed by laborers (32%), businessmen (12%), housewives (9%), farmers (7%) and landlords (4%). 33% did not have any formal education, 14% had education till primary, 7% till secondary, 31% till matriculation, 8% till intermediate and 7% were graduation and above as depicted in Table I.

Majority of people had good knowledge of the mode spread of dengue fever that is by biting of Aedes mosquito (81%) while others suggested air droplet (10%), physical contacts (6%) and uncooked food (3%) can spread dengue fever. 63% respondents claimed clean stagnant water as place of growth, 23% dirty stagnant water, 8% clean flowing water, 5% dirty flowing water and 1% had no idea of place of growth of vector. About 44% respondents suggested dengue spread was common in months of June to August while others suggested March to May (35%), September to November (11%) and December-February (10%) were common seasons for spread. Biting time of Aedes mosquitoes according to respondents was early morning (35%), night (28%), evening (25%) and 12 o'clock at noon (12%).

Based on the scoring of the knowledge section regarding symptoms of dengue fever most of respondents selected high grade fever(40%), rash (20%), bleeding(11%), pain(3%), all of above(16%) however 10% had no idea of symptoms. Respondents reported that treatment strategy if they contacted DF will be allopathic (74%), homeopathic (9%), hakeem (9%) and no treatment at all (8%). When asked about management for dengue fever 41% did not have any idea while others responded that blood transfusion (21%), antipyretic (17%), fluid intake (13%) and all of above (8%) were included. Regarding outcomes of dengue 58% responded that recovery is common, 33% responded death will occur, 4% lifelong progression and 5% had no idea about outcome.

With regard to the practices of preventive measures 55% had no awareness while others responded with wearing full sleeves and use of repellents (17%), covering water container (16%), insecticide spray (10%) and by not staying outdoors in evening (2%). According to respondents they considered television as better source of information (58%) followed by news paper (19%), friends and relatives (12%) internet (11%). Knowledge, attitude and practices are summarized in Table II.

TABLE I: DEMOGRAPHIC CHARACTERISTICS

Total Number of Respondents (n)	100
Gender	
Male	63%
Female	37%
Mean Age	30.24±16.57 years
Age Group	
16-30 years	68%
30-45 years	16%
40-60 years	9%
61-75 years	7%
Occupation	
Students	36%
Laborers	32%
Businessmen	12%
Housewives	9%
Farmers	7%
Landlords	4%
Education	
None	33%
Primary	14%
Secondary	7%
Matric	31%
Intermediate	8%
Graduation & Above	7%

TABLE II: KNOWLEDGE, ATTITUDE AND PRACTICES

Mode of Spread		Treating Person	
Biting of Aedes mosquito	81%	Allopathic	74%
Air droplet	10%	Homeopathic	9%
Physical contacts	6%	Hakeem	9%
Uncooked food	3%	No treatment	8%
Place of Growth		Management	
Clean Stagnant Water	63%	Blood Transfusion	21%
Dirty Stagnant Water	23%	Antipyretic	17%
Clean Flowing Water	8%	Fluid intake	13%
Dirty Flowing Water	5%	All of above	8%
No Idea	1%	No Idea	41%
Season of Spread		Outcome of Disease	
June-August	44%	Recovery	58%
March-May	35%	Death	33%
September-November	11%	Lifelong Progression	4%
December-February	10%	No Idea	5%
Biting Time of Mosquitoes		Preventive Measures	
Early Morning	35%	No awareness	55%
Night	28%	Wearing full sleeves and use of repellents	17%

Evening	25%	Covering water container	16%
12 o'clock at Noon	12%	Insecticide Spray	10%
		Not staying outdoors in evening	2%
Symptoms Of Dengue Fever		Source Of Information	
Fever	40%	Television	58%
Rash	20%	Newspaper	19%
Bleeding	11%	Relatives and friends	12%
Retero-orbital Pain (3%)	3%	Internet	11%
All of above	16%		
No Idea	10%		

DISCUSSION

This study demonstrates that there is lack of awareness among masses regarding dengue fever, its mode of spread, its symptoms, preventive measures and outcome of disease. Majority of our respondents were young adult male having an outdoor job and lived in rural areas. This study demonstrates the similar sociodemographic features as patients of dengue fever as depicted by Afzal M 2014¹² in their study done in Gujrat. They showed that dengue fever is more prevalent in rural areas, in low socioeconomic groups, having potential breeding places around their homes and those having at outdoor nature of job. Respondents showed a good knowledge regarding vector of dengue fever (81%) which is comparable to other studies conducted in Pakistan^{7,8,13} and clean stagnant water as its breeding sites(63%). Stagnant water is major cause of spread of dengue fever in Pakistan¹⁴ as depicted by one study conducted in Faisalabad and in another study conducted in Lahore¹⁵. Respondents had poor knowledge regarding season of spread of dengue fever and only 11% reported that its common in September to November contrary to study conducted by Suleman M et al. who concluded that maximum number of confirmed cases are reported from September to November¹⁶ but they did have a good knowledge of biting time of mosquitoes that is day time which is also concluded by other studies^{7,8,13}. Only 10% of respondents had no idea about the symptoms of dengue fever while others had a considerably good knowledge of symptoms and recognized fever as most common symptoms. Adequate knowledge of symptoms has been depicted among common population in other studies from Pakistan^{8,13}. Most of respondents showed that they will approach a physicians in case of dengue fever but 26% respondents had alternative options like hakeem, homeopathic or no treatment which quite alarming as it may increase mortality among patients and many cases will go unidentified. This relates to knowledge about management of dengue fever as most of them

had no idea about it although some of them did identify anti-pyretics as treatment option showing similar results to other studies in Pakistan¹³. Most of respondents perceived that recovery is possible from dengue fever(58%) while others recognized dengue as deadly disease although knowledge of outcome is less as compared to another study⁷.

Respondents had poor awareness regarding preventive measures of dengue fever. 55% had no awareness regarding any preventive measure which is contrary to studies by Qadir S 2015⁷ and Itrat A et al¹³ but coincides with results of Syed M et al.⁵ which showed that awareness is less among low socioeconomic groups. Awareness regarding prevention was also very poor as compared to studies in other population groups^{7,13}. Electronic media was reported as most effective source of awareness by our study population which is reported by other studies in a similar way^{7,17} but contrary to study by Bota R 2014⁸ which shows relatives and friends as most common sources of awareness.

This study shows that rural population has overall poor knowledge, attitudes and practices regarding dengue fever and there is need to further enhance the awareness programs by governments as they have proven to less effective^{4,18}. Other studies in Pakistan have shown similar results^{4,7-9,13,18}. There is need for integrated community approach via social, political mediums and through electronic media and the areas endemic should be focused in awareness programs. Emergency facilities along with medicines, mosquito repellents, chemical and mosquito nets should be provided free of cost or with controlled enhancing their use and thus dengue fever prevention.

CONCLUSION

The rural population of Gujrat has good knowledge regarding mode of spread, symptoms of dengue fever and where to get treatment however they lacked knowledge in management, outcome and prevention of dengue fever.

ACKNOWLEDGEMENT

We are thankful to Maria Mehmood, Narisa Pervez and Sania Riaz final year medical students at Nawaz Sharif Medical College, Gujrat for their help in data collection.

REFERENCES

1. World Health Organization. Dengue Fever. Available at <http://www.emro.who.int/pak/programmes/dengue-fever.html>.
2. National Institute of Health. MNHS-R&C. Govt. of Pakistan. Seasonal awareness and alert letter (SAAL) in 36th issue. June–September 2016. Available at <https://www.nih.org.pk/wp-content/uploads/2018/03/Seasoanl-Awareness-and-Alert-Letter-SAAL-36th-Issue.pdf>
3. Haider Z, Ahmad FZ, Mahmood A, Waseem T, Shafiq I, Raza T, et al. Dengue fever in Pakistan: a paradigm shift; changing epidemiology and clinical patterns. *Perspect Public Health* 2015;135(6): 294-8. doi: 10.1177/1757913915599019.
4. Zahir A, Ullah A, Shah M, Mussawar A. Barriers to prevention in dengue fever in Pakistan, Khyber Pakhtunkhwa, Swat district. *JADE* 2016; 5(6):73-82. Doi. 10.13014/k28p5xpg.
5. Syed M, Saleem T, Syeda UR, Habib M, Zahid R, Bashir A, et al. Knowledge, attitudes and practices regarding dengue fever among adults of high and low socioeconomic groups. *J Pak Med Assoc* 2010; 60(3): 243-7.
6. Lakhia JA, Sajid HMU, Khan MA, knowledge, attitude and practices relating to dengue fever among teachers of Peshawar Garrison. *Pak J Physiol* 2016; 12(2): 40-2.
7. Qadir S, Ahmad I, Akhtar MN, Naeem H. Knowledge, attitude and practice about dengue fever among local population. *Gomal J Med Sci* 2015; 13(2): 87-90.
8. Bota R, Ahmed M, Jamali MS, Aziz A. Knowledge, attitude and perception regarding dengue fever among university students of interior Sindh. *J Infect Public Health* 2014; 7(3):218–23. doi: 10.1016/j.jiph.2013.11.004
9. Siddiqui TR, Ghazal S, Bibi S, Ahmed W, Sajjad SF, Stuyft P Van der. Use of the health belief model for the assessment of public knowledge and household preventive practices in Karachi, Pakistan, a dengue-endemic city. *PLoS Negl Trop Dis* 2016; 10(11): e0005129. doi: 10.1371/journal.pntd.0005129
10. Pakistan Bureau of Statistics. Population Density and Urban Rural Proportion. Available at <http://www.pbs.gov.pk/content/area-population-density-and-urbanrural-proportion>.
11. Rehman A, Jingdong L, Hussain I. The province-wise literacy rate in Pakistan and its impact on the economy. *Pac Sci Rev B: Humanities Soc Sci* 2015; 1:140-44.
12. Afzal M, Tarar SH, Akhtar M. Socio-demographic Characteristics and Clinical Spectrum of Dengue patients presenting to Aziz Bhatti Shaheed Teaching Hospital, Gujrat. *Pak J Med Health Sci* 2014; 8(1): 56-60
13. Itrat A, Khan A, Javaid S, Kamal M, Khan H, Javed S, et al. Knowledge, awareness and practices regarding dengue fever among the adult population of dengue hit cosmopolitan. *PLoS One* 2008; 3(7):e2620. <https://doi.org/10.1371/>

- journal.pone.0002620
14. Muhammad I, Khan J, Iqbal H, Rahman H, Muhammad A, Seroprevalence and epidemiological status of dengue viral infection in remote areas of Pakistan. *Asian Pac J Trop Med* 2016; 6(10): 776-7. [https://doi.org/10.1016/S2222-1808\(16\)61128-3](https://doi.org/10.1016/S2222-1808(16)61128-3)
 15. Maqbool A, Mushtaq M, Ahmed MUD, Nisar M. Study of socio-demographic risk factors of dengue fever in a tertiary care hospital in Pakistan. *Int J Infect Dis* 2014; 21(1):257-8. <https://doi.org/10.1016/j.ijid.2014.03.956>.
 16. Suleman M, Lee HW, Zaidi SS, Alam MM, Nisar N, Aamir UB, et al. Preliminary Seroepidemiological survey of dengue infections in Pakistan, 2009–2014. *Infect Dis Poverty* 2017; 6:48. doi: 10.1186/s40249-017-0258-6.
 17. Ur Rehman A, Mahmood MA, Kazmi SF, Munir F, Ghani U. Dengue Fever; Impact of Knowledge on Preventive Practice. *Ann. Pak. Inst. Med. Sci* 2015;11(4):195-201.
 18. Saeed Q, Ali B, Jaleel W, Naqqash MN, Sial MU, Ghouri F, et al. Effectiveness of dengue fever eradication program in Southern Punjab, Pakistan: A cross-sectional survey. *J. Entomol. Zool. Stud* 2017; 5(2): 125-9.



AUTHOR AFFILIATION:

Dr. Syed Muhammad Ali Shah (*Corresponding Author*)

Department of Medicine
Aziz Bhatti Shaheed Teaching Hospital
Gujrat, Punjab-Pakistan.
Email: syedmuhammadalishah5@gmail.com

Dr. Kanza Malik

WMO, Department of Medicine
Aziz Bhatti Shaheed Teaching Hospital
Gujrat, Punjab-Pakistan.

Dr. Ali Malik

PGR, Department of Surgery
DHQ Hospital, Rawalpindi-Pakistan.

Dr. Maryam Abid

WMO, Sheikh Zaid Hospital
Rahim Yar Khan, Punjab-Pakistan.

Dr. Sofia Bano

WMO, National Hospital
Lahore, Punjab-Pakistan.